



Effects of Walking Football During Ramadan Fasting on Heart Rate Variability and Physical Fitness in Healthy Middle-Aged Males.

Nouredine Kammoun, Sami Hidouri, Amine Ghram, Achraf Ammar, Liwa Masmoudi, Tarak Driss, Beat Knechtle, Katja Weiss, Omar Hammouda, Mehdi Chlif

► To cite this version:

Nouredine Kammoun, Sami Hidouri, Amine Ghram, Achraf Ammar, Liwa Masmoudi, et al.. Effects of Walking Football During Ramadan Fasting on Heart Rate Variability and Physical Fitness in Healthy Middle-Aged Males.. American journal of men's health, 2022, 16 (3), pp.15579883221103418. 10.1177/15579883221103418 . hal-03703791

HAL Id: hal-03703791


<https://u-picardie.hal.science/hal-03703791>




Submitted on 24 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of Walking Football During Ramadan Fasting on Heart Rate Variability and Physical Fitness in Healthy Middle-Aged Males

American Journal of Men's Health
May-June 1–15
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15579883221103418
journals.sagepub.com/home/jmh


Noureddine Kammoun¹ , Sami Hidouri^{1†}, Amine Ghram^{2,3†},
Achraf Ammar^{4,5}, Liwa Masmoudi¹, Tarak Driss³, Beat Knechtle⁶ ,
Katja Weiss⁷, Omar Hammouda^{3,8†} , and Mehdi Chlif^{9,10†}

Abstract

This study aimed to investigate the effect of a walking football (WF) program during Ramadan fasting (RF) on heart rate variability (HRV) indices, body composition, and physical fitness in middle-aged males. Thirty-one healthy sedentary men were randomized to WF ($n = 18$) and control ($n = 13$) groups. Both groups participated in RF. The WF group were involved in a training program (small-sided games) of three sessions a week during RF. The time and frequency domains of HRV, body composition, handgrip, lumbar strength, Modified Agility Test (MAT), and 6-minute walk test (6MWT) were measured before Ramadan (BR), during Ramadan (DR), and after Ramadan (AR). We reported that RF has significantly altered some parameters of HRV DR; the mean HR decreased while the mean RR, LF, and HF increased. WF had a significant effect on HRV and mean HR DR compared with BR and AR decreased while mean RR, HF and LF increased. DR, body mass decreased in both groups, while body mass index (BMI) decreased and lean mass increased only in WF group. Lower body mass and BMI levels were reported AR only in WF group. Physical capacity improved AR, compared with BR, only in the WF group with longer distance in 6MWT, shorter time(s) in MAT, and higher lumbar strength levels. We conclude that RF increases parasympathetic system activity. WF practice during RF is safe and might improve body composition, physical fitness, autonomic cardiac function, and physical fitness in middle-aged males.

Keywords

autonomic heart function, walking football, Ramadan, health, fitness

Received March 11, 2022; revised May 7, 2022; accepted May 10, 2022

Introduction

Holy Ramadan is the ninth lunar month (29–30 days long). Fasting is one of the five pillars of Islam (Adawi et al., 2017). Healthy Muslims fast from dawn to dusk (12–20 hr/day, depending on the season and the geographical region). Observers refrain from eating food, drinking liquids, and other physical needs such as smoking and sexual intercourse (Ghram et al., 2021; Jahromi et al., 2014). Ramadan fasting (RF) as intermittent fasting is mandatory for all healthy adult Muslims (Akbari et al., 2022; Lessan & Ali, 2019). Several studies have been conducted over the last decade to investigate the impact of RF on physiological performance (Maughan et al.,

2010; Zerguini et al., 2007). Because of its potential positive effects on health and aging, intermittent fasting is linked to enhanced longevity (Persynaki et al., 2017). Importantly, fasting has been associated with some heart benefits (Nematy et al., 2012) including decreased cardiac events (Temizhan et al., 1999).

Heart rate variability (HRV) can reflect the adjustment of heart rate (HR) by providing a picture of the dynamic balance between sympathetic and parasympathetic branches of the autonomic nervous system (ANS) (Aubert et al., 2003; White & Raven, 2014). HRV serves, therefore, as an important index of pathological changes and is widely used to diagnose cardiovascular risk (Ammar et al., 2021; Kubota et al., 2017). Decreased HRV is



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

associated with advancing age, increased risk of cardiac events in clinically disease-free patients (Tsuiji et al., 1996), and higher mortality rates (Kleiger et al., 1987; Odemuyiwa et al., 1991). Increased HRV is associated with lower mortality rates (Sandrone et al., 1994; Yusuf et al., 1985). HRV is reduced during periods of stress and can be increased with exercise (Prinsloo et al., 2014). Indeed, it was previously reported that athletes have a higher HRV than sedentary individuals (Goldsmith et al., 1992).

Although widely credited as an effective marker for cardiovascular health and as a noninvasive tool to estimate autonomic activity, limited number of studies investigated the effects of fasting on HRV, with no clear consensus. While Mazurak et al. (2013), reported that an acute (48 h) total fast induced a decrease in total HRV under baseline conditions, Rodrigues et al. (2019), reported no significant alteration in HRV following an acute fasting of 12 hr in healthy adults. To the best of the authors' knowledge, only one study has examined the chronic effect of chronic intermittent fasting and reported higher HRV in the fasting group compared with the non-fasting group after Ramadan (Cansel et al., 2014). Thus, further studies are needed to confirm whether RF is a sufficient stimulus to improve HRV in healthy adults.

Football practice is beneficial to cardiorespiratory, musculoskeletal, and metabolic systems (Krustrup et al., 2010). Recreational football might help restore the fitness and cardiac function of aging sedentary adults (Andersen et al., 2014; Schmidt et al., 2014) and maintain an active lifestyle (Luo et al., 2018). Walking football (WF) has been recently recognized as a variant of football and credited as an optimal, motivating, and enjoyable activity for older adults (Cholerton et al., 2021; Madsen et al., 2021; Reddy et al., 2017; Zainudin et al., 2021). Twelve weeks

of WF reduce heart disease and stroke risks and improve blood pressure (BP) of people aged 50 to 65 years (Reddy et al., 2017). It was demonstrated that HRV remained unchanged after low-to-moderate aerobic exercise in fasting and nonfasting healthy individuals (Rodrigues et al., 2019) and after moderate-intensity exercise with low and moderate carbohydrates intake (Lima-Silva et al., 2010).

Fasting and exercise training are used as nonpharmacological interventions to investigate the regulation of intermediary metabolism (Maughan et al., 2010). Fasted-state training may decrease body weight, fat-free mass, and fat mass (Zouhal et al., 2020). Recent studies recommend WF for fasting Muslims. (Zainudin et al., 2021). Further studies are needed to determine the concomitant effects of RF and exercise on factors affecting both the ANS and physical capacity. There is also little information about the impact of RF only on HRV parameters. This study aimed to investigate the effect of a WF program during RF on HRV indices, body composition, and physical fitness in middle-aged males. We hypothesized that WF would improve HRV, body composition, and fitness parameters in middle-aged adults.

Materials and Methods

Ethical Approval

The study was approved by the local research ethics committee (*information anonymized for peer review*) and the protocol was conducted according to the Declaration of Helsinki. Participants received a detailed description of the study protocol, including the possible risks and benefits associated with the investigation; and submitted written informed consent before participation.

¹Research Laboratory Education, Motricité, Sport et Santé (EM2S) LR19JS01, High Institute of Sport and Physical Education of Sfax, University of Sfax, Sfax, Tunisia

²Healthy Living for Pandemic Event Protection (HI—Pivot) Network, Chicago, IL, USA

³Department of Exercise Physiology, Faculty of Physical Education and Sport Sciences, University of Tehran, Tehran, Iran

⁴Interdisciplinary Laboratory in Neurosciences, Physiology and Psychology: Physical Activity, Health and Learning (LINP2), UFR STAPS, UPL, Paris Nanterre University, Nanterre, France

⁵Department of Training and Movement Science, Institute of Sport Science, Johannes Gutenberg University Mainz, Mainz, Germany

⁶Institute of Primary Care, University of Zurich, Zurich, Switzerland

⁷Medbase St. Gallen Am Vadianplatz, St. Gallen, Switzerland

⁸Research Laboratory, Molecular Bases of Human Pathology, LR19ES13, Faculty of Medicine, University of Sfax, Sfax, Tunisia

⁹EA 3300 "APS and Motor Patterns: Adaptations-Rehabilitation," Sport Science Department, Picardie Jules Verne University, Amiens, France

¹⁰Tunisian Research Laboratory Sports Performance Optimization, National Center of Medicine and Science in Sports (NCMSS), Tunis, Tunisia

†These authors contributed equally to this work as second authors.

‡These authors contributed equally to this work as last authors.

*Tarak Driss and Omar Hammouda are now affiliated to Interdisciplinary Laboratory in Neurosciences, Physiology and Psychology: Physical Activity, Health and Learning (LINP2), UFR STAPS, UPL, Paris Nanterre University, Nanterre, France

Corresponding Author:

Mehdi Chlif, Tunisian Research Laboratory Sports Performance Optimization, National Center of Medicine and Science in Sports (NCMSS), Ave Med Ali Akid, El Menzah, Tunis 263, Tunisia.

Email: mehdi.chlif@gmail.com




Figure 1. The Consort Flow Diagram.

Participants

A total of 44 participants were initially enrolled in this study between May and June 2019. The inclusion criteria were age 45 years or older, healthy, willing to fast throughout Ramadan, and the ability to provide informed consent. All participants were nonsmokers. Participants with orthopedic, cardiovascular, pulmonary, neurologic or metabolic issues were excluded.

According to the International Physical Activity Questionnaire classification, healthy inactive, irregularly active, and active people were included (Craig et al., 2003). Participants classified as “very active” were considered not representative of the general population (outliers) and were excluded. Of the 44 participants, four declined to participate in the study, and four did not meet the inclusion criteria. Finally, 36 participants (body mass index [BMI] < 30) were included in the study and were randomly assigned to the WF group ($n = 18$) who participated in RF and WF training or the control group (CON, $n = 18$) who participated only in RF and were not receive any physical activity interventions (Figure 1).

In WF, we retained the data of subjects who participated in at least 90% of the training schedule.

Randomization and Blinding

The randomization sequence (ratio 1:1) was generated using a computerized block randomization generator

(randomizer.org, accessed on April 20, 2019). An independent study coordinator conducted the allocation sequence, who was not involved in the assessment and data analysis processes, enrolling participants in sequentially numbered, opaque, sealed envelopes.

Experimental Design

The study was conducted in Sfax, Tunisia, in 2019. Ramadan lasted from May 6 to June 5. The length of each daytime fast was approximately 15 to 16 hr. During the fasting period, participants abstained from food, drink, smoking, and sexual activity between dawn and sunset. The participants were familiarized with testing procedures and measurements before the start of the study.

To reduce the impacts of circadian rhythms, all testing sessions were held in the morning at the same time of day and in the same order. Fitness assessment included anthropometric measurements and assessment of cardio-respiratory capacity, muscular strength and flexibility through field tests. HRV data were collected in a silent room, with controlled temperature and relative humidity. Tests were performed in three separate sessions. The first session was performed one week prior to Ramadan (BR) and the second and the third sessions at the end of the second week (DR) and 7–10 days after Ramadan (AR), respectively (Figure 2).

During Ramadan, WF participated in a 4-week training program with three sessions per week interspersed by




Figure 2. Experimental Design.

at least 48 hr of recovery. Training sessions were scheduled from 4:00 pm to 6:00 pm. Each session lasted ~65 min: 10 min warm-up, two 20-min periods of play with a half time of 10 min, 5 min of stretching. During the 20-min period of play, WF played small-sided games 4 versus 4 or 5 versus 5 on a football pitch of 30 to 45 m \times 45 to 60 m, respectively. Meanwhile, CON remained inactive, as they were instructed to maintain their current lifestyle until the end of the study and simply underwent the assessment tests.

HRV Analysis

HRV indices were recorded using Polar technology (Polar V800, Polar Electro Oy, Kempele, Finland). This device has been considered a reliable and valid method of capturing and analyzing interbeat variability (Giles et al., 2016). Data were collected between 10 am and 11 am to avoid a possible circadian influence on the autonomic function. Before the test, participants rested comfortably in a supine position for at least 10 min in a quiet dimly lit room with constant temperature. During the test, the participants were asked to lie still, close their eyes, abstain from talking, and maintain normal breathing frequency. Tidal volume was maintained during the testing at a constant breathing rate of 0.25 Hz by listening to recorded breathing sounds at this rate to neutralize the effect of respiratory sinus arrhythmia on HRV. The test duration gives a good definition of frequency-domain measures and an acceptable resolution of time-domain measures ("Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the

European Society of Cardiology and the North American Society of Pacing and Electrophysiology," 1996; Shaffer et al., 2020). Before signal analysis, all ectopic beats were filtered and corrected. To analyze the HRV, we calculated the following parameters in the time domain: the mean of the RR intervals, the standard deviation of the normal RR interval (SDNN), the root mean square of the successive differences of the R-R intervals (RMSSD), and the percentage of successive interval differences larger than 50 ms (PNN50). The frequency-domain parameters were obtained based on the fast Fourier transformation to quantify the low-frequency bands (LF) (0.04–0.15 Hz), high frequency (HF) (0.15–0.40 Hz), and the ratio LF/HF expressed in units (n.u) were measured. In the nonlinear domain, we used SD1 as an index of the instantaneous variability of beat-to-beat data. Data analysis was performed by the same researcher using the standard Kubios HRV software (v. 3.1.1.).

Body Composition

Body mass was measured to the nearest 0.1 kg on a digital scale (Tanita TBF 401, Tanita Corp., Japan), and height was measured to the nearest 0.1 cm on a stadiometer (Holtain, Crymych, Dyfed, UK). BMI was computed by dividing body mass in kilograms by height in square meter (kg/m^2). A foot-to-foot bioelectric impedance analyzer was used to determine body fat percentage (BF%), fat-free mass (FFM), and total body water (TBW) content (Tanita TBF 401, Tanita Corp., Japan). This method has been validated against the reference methods (Bosy-Westphal et al., 2008). The bioelectrical




Figure 3. Modified Agility T-Test (MAT).

impedance measurements were carried out following the manufacturer's recommendations. For the measurement, subjects were requested to remove all clothing (excluding shorts), shoes, jewelry, and other accessories. Gender, height, and level of physical activity were manually entered into the keypad interface. Subjects were measured while standing erect with their bare feet on the footpads of the analyzer.

6-Min Walk Test

The 6-min walk test (6MWT) was performed outdoors along a flat, straight, 30-m walking course and measured using the better of two tests separated by ≥ 30 min as per ATS guidelines (Crapo et al., 2002). HR was recorded before and after the 6MWT. Participants were encouraged every minute during the test. They were allowed to stop and rest during the test but were instructed to resume walking as soon as they felt able to do so. We used a stopwatch to measure test time and a HR monitor (Polar V800, Polar Electro Oy, Kempele, Finland) to assess HR. At the end of the test, the 6-min walking distance (6MWD) was recorded.

Modified Agility T-Test

The Modified Agility T-Test (MAT) was used to assess the speed of participants while measuring their agility by using photoelectric cells (Figure 3). The participants performed the MAT by completing multidirectional sprinting, shuffling, and backpedaling (Hickey et al., 2009; Sassi et al., 2009; Scanlan et al., 2019). Each participant was required to perform three trials separated by 5 min of rest for recovery. The participants started at cone A, made a forward linear sprint of 5 m to cone B, a leftward shuffle of 2.5 m to cone C, a rightward shuffle of 5 m to cone D, a leftward shuffle of 2.5 m back to cone B, and finally a linear backpedal of 5 m to cone A.

Hand Grip Strength

The maximum strength of the forearm and hand muscles assessment was conducted using a hydraulic dynamometer (Model SH5001, Saehan Corporation, Korea) following the American Society of Hand Therapists recommendations (MacDermid et al., 2015). Subjects were seated with the arms parallel to the body, flexed at the elbow to 90°, and the forearm in a neutral position. Wrist flexion or extension was not allowed. Each participant repeated the test three times and only the highest value was considered. The dominant side was defined according to the self-reported preferred hand while performing daily activities (Günther et al., 2008). The subjects were instructed to perform their maximum isometric effort for 5 s during the test with a 1-min rest interval between the tests. The standardized instructions “When I say go, grip as hard as you can until I say stop” and “Harder . . . harder . . . harder . . . relax” were given prior and throughout the test, respectively (Innes, 1999; MacDermid et al., 2015). No other body movements were allowed.

Lumbar Strength

A calibrated lumbar strength dynamometer (BASELINE, New York, USA) was used to measure the static strength of back muscles. The dial ranges from 0 to 300 kg. Participants stand on the platform with their legs shoulder-width apart, knees extended, arms outstretched, chest bent toward the hips and palms facing down. They had to bow down enough to reach the dynamometer chain, which should be perpendicular to the platform. The experimenter faced the participant, offset to the side, and held the participant's hips to prevent any movement. After the demonstration and a familiarization trial, three trials were performed. Then, participants pulled the chain until their back was perfectly straight with rest periods of 2 min between trials. Maximal strength for the three trials was used for further analysis.

Statistical Analysis

Data analyses were performed using SPSS version 25 for Windows (SPSS Inc, Chicago, IL, USA). Values are presented as means \pm standard deviation (SD). The Shapiro–Wilk test was performed to check the normality of data. Two-way ANOVA with repeated measures was used to determine the differences between groups (2-Condition Group: WF or CON \times 3 Times of Measurement: BR, DR, AR. When a difference was identified, a Bonferroni post hoc test was used. An a priori power analysis was conducted using G*Power3 (Faul et al., 2007) to test the difference between two independent group means using a

two-tailed test, a medium effect size ($d = .50$), and an alpha of .05. Results reported that a total sample of 44 participants was required to achieve a power of .95. Effect sizes were calculated as partial eta-squared (η^2p) for the ANOVA to estimate the meaningfulness of significant findings. η^2p values of 0.01, 0.06, and 0.13 represent small, moderate, and large effect sizes, respectively. Statistical significance was assigned at $p < .05$ for all analyses.

Results

Population Characteristics

Of the 36 study participants, 5 participants in the CON opted out of the study: personal reasons ($n = 1$), back pain ($n = 1$), unexpected work ($n = 2$), moving to other country ($n = 1$). Therefore, 31 participants completed the study (18 in the WF and 13 in the CON) (Figure 1). Baseline characteristics of participants are reported in Table 1. There were no significant differences in baseline characteristics WF and CON, indicating the homogeneity of the two groups.

HRV

HRV is reported in Table 2. Two-way ANOVA with repeated measurements of mean HR and mean RR data showed a significant interaction (Ramadan \times Group). The statistical analysis reported a significant main effect of Ramadan on Mean HR, mean RR, HF, and LF. Mean HR was significantly lower in DR compared with BR ($p = .000$) and in AR ($p = .002$) in WF. Mean HR was significantly lower AR in WF compared with CON ($p = .010$).

Mean RR was significantly higher in DR compared with BR ($p = .000$) and in AR compared with DR AR ($p = .000$) in WF. Mean RR was significantly lower AR compared with DR in CON ($p = .018$) and was significantly higher in WF compared with the CON ($p = .001$). In addition, HF and LF were significantly higher DR compared with BR ($p = .022$ for HF and LF) and in AR compared with DR ($p = .026$ for HF; and $p = .017$ for LF) in WF. A significant main effect of the group was detected only in LF/HF ($p = .022$). No significant changes were detected for SDNN, RMSSD, PNN%, SD1, and LF/HF.

Body Composition

Body composition is reported in Table 3. The statistical analysis reported a significant main effect of Ramadan on body mass, BMI, fat mass (%), lean mass (%), and TBW (kg). Body mass was significantly lower DR ($p = .000$)

Table 1. Baseline Anthropometric Data Characteristics.

	WF ($n = 18$)	CON ($n = 13$)
Age (year)	53.70 \pm 7.3	52.75 \pm 4.2
Body mass (kg)	89.1 \pm 9.7	84.6 \pm 9.7
Height (cm)	177.75 \pm 5.1	175.46 \pm 5.8
BMI (kg/m ²)	28.2 \pm 3	27.9 \pm 2.3
Fat mass (%)	23.7 \pm 5.3	23.4 \pm 5.6
Lean mass (%)	54.2 \pm 4	55.2 \pm 3.6

Values are presented as mean \pm SD.

WF = walking football group; CON = control group; BMI = body mass index.

and AR ($p = .000$) compared with BR in WF and DR compared with BR in CON ($p = .004$). BMI was significantly lower DR ($p = .000$) and AR ($p = .005$) compared with BR in WF. Lean mass increased only in the WF group DR compared with BR ($p = .005$) while AR it increased for both groups ($p = .001$ for WF; and $p = .004$ for CON). Fat mass (%) was significantly lower AR compared with BR in CON ($p = .016$). However, there is no significant effect on TBW in both groups.

6MWT and MAT

The results of 6MWT, MAT, and strength performance are reported in Table 4. The two-way ANOVA with repeated measurements of 6MWD and MAT reported a significant interaction (Ramadan \times Group; $p = .000$; and $p = .018$, respectively). The statistical analysis reported a significant main effect of Ramadan on 6MWD, HRmax, mean HR, and MAT ($p = .000$) and the main effect of group on 6MWD ($p = .006$). 6MWD was significantly lower DR and AR in CON compared with WF ($p = .000$). 6MWD was significantly higher DR ($p = .036$) and AR ($p = .000$) compared with BR in the WF. HRmax was significantly higher AR compared with DR only in WF ($p = .023$). Also, HRmax was significantly higher DR in WF compared with CON ($p = .008$). MAT performance was significantly lower AR compared with BR ($p = .000$) and DR ($p = .000$) in WF. MAT was significantly lower DR ($p = .000$) and AR ($p = .013$) in WF compared with CON.

Handgrip and Lumbar Strength

Handgrip and lumbar strength are reported in Table 4. Statistical analysis reported significant interaction (Ramadan \times Group) only on lumbar strength ($p = .027$) and a nonsignificant main effect of Ramadan and main effect of group on Gripping force and Lumbar strength. Lumbar strength was significantly higher after R compared with before R ($p = .019$) in WF group.

Table 3. Body Composition, Before, During, and After Ramadan for Both Groups.

Parameters		BR	DR	AR	Group effect			Ramadan effect			Group \times Ramadan		
					$F_{(1, 29)}$	p value	$\eta^2 p$	$F_{(2, 58)}$	p value	$\eta^2 p$	$F_{(2, 58)}$	p value	$\eta^2 p$
Body mass (kg)	CON	84.6 \pm 9.7	82.8 \pm 9.5 ^{aa}	83.4 \pm 9.7	1.019	0.321	0.034	39.452	0.000	0.576	4.497	0.015	0.134
	WF	89.1 \pm 9.7	85.8 \pm 9.4 ^{aaa}	86.4 \pm 9.3 ^{aaa}									
BMI (kg/m ²)	CON	27.9 \pm 2.3	27.4 \pm 2.3	27.5 \pm 2.5	0.002	0.965	0.000	17.453	0.000	0.376	2.599	0.083	0.082
	WF	28.2 \pm 3	27.1 \pm 2.8 ^{aaa}	27.5 \pm 2.8 ^{aa}									
Fat mass (%)	CON	23.4 \pm 5.6	21.9 \pm 3.1	20.7 \pm 3.6 ^a	0.048	0.828	0.002	11.018	0.000	0.275	0.500	0.609	0.017
	WF	23.7 \pm 5.3	21.8 \pm 5.3	21.6 \pm 5.1									
Lean mass (%)	CON	55.2 \pm 3.6	56 \pm 2.3	57 \pm 2.7 ^{aa}	0.391	0.537	0.013	16.309	0.000	0.360	1.050	0.357	0.035
	WF	54.2 \pm 4	55.8 \pm 4.1 ^{aa}	55.9 \pm 3.8 ^{aa}									
TBW (kg)	CON	62.2 \pm 8.5	62.1 \pm 7	63.4 \pm 6.9	0.651	0.426	0.022	3.399	0.040	0.105	0.118	0.889	0.004
	WF	64.3 \pm 6.6	64.1 \pm 5.4	65 \pm 6.4									

Values are means \pm SD; BMI = body mass index; TBW = total body water; BR = before Ramadan; DR = during Ramadan; AR = after Ramadan; WF group = walking football group.

^aSignificantly different from BR at $p < .05$; ^{aa}Significantly different from BR at $p < .01$; ^{aaa}Significantly different from BR at $p < .001$.

Table 4. Effect of Ramadan Fasting on 6MWT, Agility and Strength Test Performance Before, During, and After Ramadan for Both Groups.

Parameters		Group effect					Ramadan effect			Group × Ramadan				
		F(1,29)		p value		η ² p	F(2,58)		p value	η ² p	F(2, 58)		p value	η ² p
6MWT														
Distance (m)	CON	660 ± 56.3	662.4 ± 52.6	689.3 ± 47.7	8.786	0.006	0.233	28.579	0.000	0.496	9.502	0.000	0.247	
	WF	675.8 ± 58.8	715.7 ± 65.9 ^a	789.8 ± 73.1 ^{*ab}										
Mean HR (beats/min)	CON	109 ± 16	113 ± 17	130 ± 11	0.857	0.362	0.029	9.054	0.000	0.238	0.133	0.876	0.005	
	WF	114 ± 17	118 ± 27	131 ± 16										
Max HR (beats/min)	CON	133 ± 19	129 ± 21	148 ± 14	1.590	0.217	0.052	10.645	0.000	0.269	0.061	0.941	0.002	
	WF	138 ± 19	137 ± 25 [*]	155 ± 20 ^b										
Agility and strength test														
MAT, time (s)	CON	7.89 ± 0.75	8.33 ± 0.7	8.03 ± 0.94	1.68	0.206	0.055	8.22	0.000	0.221	4.31	0.018	0.13	
	WF	7.94 ± 0.55	8.04 ± 0.75 [*]	7.38 ± 0.7 ^{*ab}										
Lumbar strength (kg)	CON	134.3 ± 33.1	129.6 ± 31.3	132.5 ± 30.2	0.060	0.809	0.002	2.405	0.099	0.077	3.858	0.027	0.117	
	WF	125.9 ± 17.3	129.5 ± 19.5	134.7 ± 18.1 ^a										
Hand grip (kg)	CON	49.3 ± 6.6	48.3 ± 7.1	49 ± 6	1.544	0.224	0.051	1.637	0.203	0.053	1.632	0.204	0.053	
	WF	45.3 ± 6.5	45.9 ± 5.5	47.7 ± 5.1										

Values are mean \pm SD; HR: heart rate, 6MWD: 6-min walking test, MAT: modified T-test; BR = before Ramadan; DR = during Ramadan; AR = after Ramadan; WF group = walking football group; CON = control group.

^aSignificantly different from control group at $p < .05$; ^bSignificantly different from BR at $p < .05$; ^{ab}Significantly different from DR at $p < .05$.

Discussion

The objective of this study was to investigate the effects of Ramadan intermittent fasting on HRV and, the effects of WF training and RF on HRV indices, body composition, and physical fitness in middle-aged adults. This study found a significant effect of practicing WF while fasting on mean RR ($p = .029$) and HR ($p = .019$) associated with an improvement in body mass ($p = .015$), 6MWT ($p = .000$), MAT ($p = .018$) and lumbar strength ($p = .027$) performance. In addition, RF had a significant effect on four parameters of HRV DR; the mean HR decreased ($p = .000$) while the mean RR ($p = .000$), LF ($p = .020$) and HF ($p = .012$) increased. In addition, RF had a significant effect on body composition; body mass ($p = .000$), lean mass ($p = .000$), BMI ($p = .000$), fat mass ($p = .000$), and TBW (Kg) ($p = .040$). Finally, a significant group effect was founded only on LF/HF ratio ($p = .022$) and on the performance 6MWT ($p = .006$).

The effect of exercise training and RF on HRV, a reflection of the cardiac ANS and a promising method for monitoring cardiovascular health, has received little attention. Our results demonstrated that RF had a minor effect on overall cardiac ANS. HRV was slightly affected by WF training indicating no significant change in the cardiac ANS activity. Therefore, our finding suggests a slight cardiac adaptation process in response to WF while fasting.

Our results corroborate in part with existing studies (Cansel et al., 2014; Hammoud et al., 2020) on the impact of Ramadan on HRV in healthy subjects. A study conducted on 80 women reported no significant difference in HRV parameters between the first and last week of Ramadan (Hammoud et al., 2020). In this study, some changes in ANS indices (e.g., HR, SDNN, RMSSD, PNN50) were observed during the fasting days between noon and shortly before the sunset [the end of the fasting period], indicating that there was a difference in ANS activity between early and late fasting (Hammoud et al., 2020). No significant difference was identified in the HRV indices between fasting and nonfasting times, except for a significantly decreased HR during fasting time (Hammoud et al., 2020). However, this result contradicts the study of Cansel et al. (2014) who reported an increase in some HRV parameters (e.g., SDNNI, RMSSD, pNN50, T power, LF, LFnu, HF, and HFnu) induced by fasting compared with the post-Ramadan period, indicating a stimulation of the parasympathetic nervous system activity in these subjects. This was due to catecholamine inhibition during fasting, which resulted in a decrease in sympathetic nervous system activity in fasting healthy subjects (Cansel et al., 2014). These findings are relatively consistent with the results of the present study as an increase was reported in mean RR, LF, and HF AR and

DR compared with BR. The discrepancy between our findings and those of the aforementioned studies could be attributed to the difference in the study participants (men vs. females), and the period and the time of day of HRV measurements.

Football is an effective way to improve health (Krustrup et al., 2013). It increases cardiovascular and musculoskeletal fitness, improves postural balance and generates high aerobic activity (Krustrup et al., 2010). However, such benefits are still to be verified in WF. To date, only a few studies have investigated the effect of WF on health conditions. One study reported no effect of a weekly WF session of 1 hr over 12 weeks on either health status or cognitive performance (Reddy et al., 2017). Another study reported a decrease in fat percentage and a rise in time to exhaustion following 12 weeks of a weekly WF session of 2 hr (Arnold et al., 2015). The influence of WF in a fasted state has not been investigated particularly for HRV parameters. This study reported that WF exercising caused no significant changes in HRV parameters. HRV is rather influenced by diet, smoking, disturbed sleep patterns (Guiraud et al., 2013; James et al., 2012), regular exercise, and exercise intensity (Hedelin et al., 2000; Madsen et al., 2021).

The 4 weeks of exercise training and the intensity of WF training, based on brisk walking, are not sufficient to affect HRV indices. Moreover, the possible sleeping habit deterioration DR could account for the absence of WF affecting health outcomes in the present study. Two previous studies on the effects of WF in older adults did not report the same health-related changes induced by traditional running football. This might be due to differences in training intensity but could also be related to differences in the volume, duration, and frequency of the training sessions (Andersen et al., 2014; Arnold et al., 2015; Reddy et al., 2017). A recent study associated a significant increase in the RMSSD index with training intensity, *which* means that a slight increase in intensity improves the regulation of cardiac function (Tornberg et al., 2019). Melanson and Freedson (2001) reported a significantly higher HRV [RMSSD, SDNN] in moderate and high-intensity aged groups. Twelve months of supervised exercise [3 months of stretching and 9 months of 5 hr/week aerobic exercise at approximately 70% of VO₂max] increased total HRV in older adults (Stein et al., 1999). In addition, our participants were exercising on an empty stomach. No changes in ANS activity were reported under these circumstances.

Physical training and fasting are credited for their positive impacts on body composition and health outcomes (Zouhal et al., 2020). Fasting and aerobic exercise training are two common ways of increasing lipolysis in adipose and muscular tissue, which reduces body fat mass (Zouhal et al., 2020). In the present study, body

composition improved in both groups. Indeed, Body mass was significantly lower DR ($p = .000$) and AR ($p = .000$) compared with BR in WF and DR compared with BR in CON ($p = .004$). BMI was significantly lower DR ($p = .000$) and AR ($p = .005$) compared with BR in WF. Lean mass increased only in the WF group DR compared with BR ($p = .005$) while AR it increased for both groups ($p = .001$ for WF; and $p = .004$ for CON). Fat mass (%) was significantly lower AR compared with BR in CON ($p = .016$). WF training while fasting was revealed to improve body mass and BMI for the WF group. The results of the present study are consistent with previous reports (Arnold et al., 2015; Reddy et al., 2017). Reddy et al. (2017) reported that WF training for 12 weeks in older adults (50–65 years old) reduced visceral fat (−1.5 kg), and fat mass (−0.38%). In addition, 12 weeks of WF training increased lean mass (+2.5 kg) and decreased body mass (−1.8), fat mass (−3%), and BMI (−1) in older adults (66 years old) with various comorbidities (Arnold et al., 2015). Thus, WF training is effective for all age groups, regardless of their health status. However, changes in body composition are rather related to RF than WF training. Compared with the pre-Ramadan period, a decrease in body mass, BMI, and fat mass (%) at the end of Ramadan has been reported (Nachvak et al., 2019). The effects of Ramadan may vary according to environmental conditions such as fasting duration, temperature, and habits (Addin Akbari et al., 2021; Faris et al., 2019, 2020; Farooq et al., 2010; Ghram et al., 2021). A previous investigation of the variations in body composition in adults during Ramadan, in a hot (45°C) and dry environment reported no significant changes in body mass, BMI, lean body mass, and TBW (Al-Barha & Aljaloud, 2019). Another study reported that RF decreased body mass, BMI, and skeletal muscle mass (Nugraha et al., 2017). Still more, another study reported a decrease in water and fat mass in obese adults after RF (Sezen et al., 2016). RF improved the body composition of obese more than of lean individuals (Harder-Lauridsen et al., 2017). Geographical, social, and cultural variables between Muslim countries and groups seem to account for the variability of results between studies.

The WF training during Ramadan reported a significant improvement in the 6MWD in the WF group. This increase in physical capacity is associated with a significant increase in the maximum HR in WF group. Data about the effects of RF on the cardiorespiratory response to exercise in sedentary adults are controversial. A small reduction (Ramadan, 2002; Ramadan & Barac-Nieto, 2000) or no impact on maximal exercise capacity and walking deficiency (Ramadan, 2002; Sweileh et al., 1992) has been reported. The causes for such inconsistent outcomes are still unknown. However, factors such as sleep deprivation or fatigue during Ramadan are crucial concerns. No significant change was observed in the gripping force, while the

lumbar strength was significantly higher AR compared with BR in WF group. In line with our findings, the levels of muscle strength (grip strength) remain unchanged during Ramadan (Bouhleb et al., 2013). Our findings demonstrate a minimal change in performance measures in WF after Ramadan, except for a substantial improvement in agility in WF group. These findings are consistent with the study of Kordi et al., who argued that the higher agility fitness level in pre-Ramadan may be due to a reduction in body mass (Kordi et al., 2011).

To the best of the authors' knowledge, this is the first study to investigate the concomitant effects of RF and WF practice on HRV parameters in middle-aged adults. The findings of the present study should be interpreted with the following limitations in mind; First, the sample size was small and the fasting and training periods were relatively short, which may partly account for the absence of significant differences between groups in most of the verified parameters. Second, the fact that all participants were males, eliminates the effect of female reproductive hormone levels on HRV and physical performance. Future studies should evaluate the effects of intermittent fasting and practice in larger male and female groups and control factors such as calorie intake, sleep pattern, psychological variables, and circadian rhythm.

Conclusion

This study reported significant improvements in HRV and physical fitness parameters after 4 weeks of WF during RF in middle-aged males. Exercise training while fasting enhances physical fitness and cardiovascular autonomic balance, which might imply that WF practice in a fasted state may contribute to exercise-induced cardio-protection. This study should prompt future investigations on the significance of ANS regulation during training exercises in a fasted state.

Acknowledgments

The authors would like to thank all the participants for their outstanding cooperation to make this paper possible.

Author Contributions

Conceptualization, N.K., O.H., and M.C.; methodology, N.K., O.H.; validation, N.K., O.H.; formal analysis, N.K., S.H., A.G., L.M.; resources, S.H.; data curation, L.M.; writing—original draft preparation, N.K., S.H., and A.G.; writing—review and editing, M.C., O.H., B.K., K.W.; AA and T.D.; supervision, M.C., O.H. All authors have read and agreed to the published version of the manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Noureddine Kammoun  <https://orcid.org/0000-0002-8422-9886>

Beat Knechtle  <https://orcid.org/0000-0002-2412-9103>

Omar Hammouda  <https://orcid.org/0000-0002-5002-687X>

References

- Adawi, M., Watad, A., Brown, S., Aazza, K., Aazza, H., Zouhir, M., Sharif, K., Ghanayem, K., Farah, R., Mahagna, H., Fiordoro, S., Sukkar, S. G., Bragazzi, N. L., & Mahroum, N. (2017). Ramadan fasting exerts immunomodulatory effects: Insights from a systematic review. *Frontiers in Immunology*, 8, 1144. <https://doi.org/10.3389/fimmu.2017.01144>
- Akbari, H. A., Ghram, A., Yoosefi, M., Arena, R. J., Lavie, C., Chtourou, H., Ben Saad, H., & Chamari, K. (2021). The COVID-19 pandemic and physical activity during intermittent fasting, is it safe? A call for action. *Biology of Sport*, 38, 729–732. <https://doi.org/10.5114/biolsport.2021.108300>
- Akbari, H. A., Yoosefi, M., Pourabbas, M., Weiss, K., Knechtle, B., Vancini, R. L., Trakada, G., Ben Saad, H., Lavie, C. J., & Ghram, A. (2022). Association of Ramadan participation with psychological parameters: A cross-sectional study during the COVID-19 pandemic in Iran. *Journal of Clinical Medicine*, 11, 2346. <https://doi.org/10.3390/jcm11092346>
- Al-Barha, N. S., & Aljaloud, K. S. (2019). The effect of ramadan fasting on body composition and metabolic syndrome in apparently healthy men. *American Journal of Men's Health*, 13(1), 1557988318816925. <https://doi.org/10.1177/1557988318816925>
- Ammar, A., Boukhris, O., Halfpaap, N., Labott, B. K., Langhans, C., Herold, F., Grassler, B., Muller, P., Trabelsi, K., Chtourou, H., Zmijewski, P., Driss, T., Glenn, J. M., Muller, N. G., & Hoekelmann, A. (2021). Four weeks of detraining induced by covid-19 reverse cardiac improvements from eight weeks of fitness-dance training in older adults with mild cognitive impairment. *International Journal of Environmental Research and Public Health*, 18(11), 5930. <https://doi.org/10.3390/ijerph18115930>
- Andersen, T. R., Schmidt, J. F., Nielsen, J. J., Randers, M. B., Sundstrup, E., Jakobsen, M. D., Andersen, L. L., Suetta, C., Aagaard, P., Bangsbo, J., & Krstrup, P. (2014). Effect of football or strength training on functional ability and physical performance in untrained old men. *Scandinavian Journal of Medicine & Science in Sports*, 24(Suppl. 1), 76–85. <https://doi.org/10.1111/sms.12245>
- Arnold, J. T., Bruce-Low, S., & Sammut, L. (2015). The impact of 12 weeks walking football on health and fitness in males over 50 years of age. *BMJ Open Sport & Exercise Medicine*, 1(1). <https://doi.org/10.1136/bmjsem-2015-000048>
- Aubert, A. E., Seps, B., & Beckers, F. (2003). Heart rate variability in athletes. *Sports Medicine*, 33(12), 889–919. <https://doi.org/10.2165/00007256-200333120-00003>
- Bosy-Westphal, A., Later, W., Hitze, B., Sato, T., Kossel, E., Gluer, C. C., Heller, M., & Muller, M. J. (2008). Accuracy of bioelectrical impedance consumer devices for measurement of body composition in comparison to whole body magnetic resonance imaging and dual X-ray absorptiometry. *Obesity Facts*, 1(6), 319–324. <https://doi.org/10.1159/000176061>
- Bouhlef, H., Shephard, R. J., Gmada, N., Aouichaoui, C., Peres, G., Tabka, Z., & Bouhlef, E. (2013). Effect of ramadan observance on maximal muscular performance of trained men. *Clinical Journal of Sport Medicine*, 23(3), 222–227. <https://doi.org/10.1097/JSM.0b013e318275d213>
- Cansel, M., Taşolar, H., Yağmur, J., Ermiş, N., Açıkgöz, N., Eyyüpkoca, F., Pekdemir, H., & Ozdemir, R. (2014). The effects of Ramadan fasting on heart rate variability in healthy individuals: A prospective study. *Anatolian Journal of Cardiology*, 14(5), 413–416. <https://doi.org/10.5152/akd.2014.5108>
- Cholerton, R., Quirk, H., Breckon, J., & Butt, J. (2021). Experiences and strategies influencing older adults to continue playing walking football. *Journal of Aging and Physical Activity*, 29, 573–585. <https://doi.org/10.1123/japa.2020-0058>
- Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., & Oja, P. (2003). International physical activity questionnaire: 12-Country reliability and validity. *Medicine & Science in Sports & Exercise*, 35(8), 1381–1395. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>
- Crapo, R. O., Casaburi, R., Coates, A. L., Enright, P. L., MacIntyre, N. R., McKay, R. T., Johnson, D., Wanger, J. S., Zeballos, R. J., Bittner, V., & Mottram, C. (2002). ATS statement: Guidelines for the six-minute walk test. *American Journal of Respiratory and Critical Care Medicine*, 166(1), 111–117. <https://doi.org/10.1164/ajrccm.166.1.at1102>
- Faris, M. A. E., Jahrami, H. A., Alhayki, F. A., Alkhawaja, N. A., Ali, A. M., Aljeeb, S. H., Abdulghani, I. H., & BaHammam, A. S. (2020). Effect of diurnal fasting on sleep during Ramadan: A systematic review and meta-analysis. *Sleep and Breathing*, 24(2), 771–782. <https://doi.org/10.1007/s11325-019-01986-1>
- Faris, M. A. E., Jahrami, H. A., Alsibai, J., & Obaideen, A. A. (2019). Impact of ramadan diurnal intermittent fasting on metabolic syndrome components in healthy, non-athletic muslim people aged over 15 years: A systematic review and meta-analysis. *British Journal of Nutrition*, 123, 1–22. <https://doi.org/10.1017/s000711451900254x>
- Farooq, S., Nazar, Z., Akhtar, J., Irfan, M., Subhan, F., Ahmed, Z., Khan, E. H., & Naeem, F. (2010). Effect of fasting during Ramadan on serum lithium level and mental state in bipolar affective disorder. *International Clinical Psychopharmacology*, 25(6), 323–327. <https://doi.org/10.1097/YIC.0b013e32833d18b2>
- Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191. <https://doi.org/10.3758/bf03193146>

- Ghram, A., Ben Saad, H., Briki, W., Jiménez-Pavón, D., Mansoor, H., Moalla, W., Akbari, H. A., Ghram, A., Al-Mohannadi, A. S., Arena, R., & Lavie, C. J. (2021). Ramadan intermittent fasting, physical activity, and covid-19 pandemic in patients with chronic diseases. *American Journal of Medicine*, 134, 1189–1191. <https://doi.org/10.1016/j.amjmed.2021.04.035>
- Giles, D., Draper, N., & Neil, W. (2016). Validity of the Polar V800 heart rate monitor to measure RR intervals at rest. *European Journal of Applied Physiology*, 116(3), 563–571. <https://doi.org/10.1007/s00421-015-3303-9>
- Goldsmith, R. L., Bigger, J. T., Steinman, R. C., & Fleiss, J. L. (1992). Comparison of 24-hour parasympathetic activity in endurance-trained and untrained young men. *Journal of the American College of Cardiology*, 20(3), 552–558. [https://doi.org/https://doi.org/10.1016/0735-1097\(92\)90007-A](https://doi.org/https://doi.org/10.1016/0735-1097(92)90007-A)
- Guiraud, T., Labrunee, M., Gaucher-Cazalis, K., Despas, F., Meyer, P., Bosquet, L., Gales, C., Vaccaro, A., Bousquet, M., Galinier, M., Sénard, J. M., & Pathak, A. (2013). High-intensity interval exercise improves vagal tone and decreases arrhythmias in chronic heart failure. *Medicine & Science in Sports & Exercise*, 45(10), 1861–1867. <https://doi.org/10.1249/MSS.0b013e3182967559>
- Günther, C. M., Bürger, A., Rickert, M., Crispin, A., & Schulz, C. U. (2008). Grip strength in healthy caucasian adults: Reference values. *Journal of Hand Surgery*, 33(4), 558–565. <https://doi.org/10.1016/j.jhsa.2008.01.008>
- Hammoud, S., Mourad, R., Karam, R., Saad, I., van den Bemt, B. J. F., & Kurdi, M. (2020). Effect of Ramadan fasting on heart rate variability as a measure of cardiac stress in a Lebanese cohort. *European Journal of Clinical Nutrition*, 74(8), 1237–1239. <https://doi.org/10.1038/s41430-020-0562-2>
- Harder-Lauridsen, N. M., Rosenberg, A., Benatti, F. B., Damm, J. A., Thomsen, C., Mortensen, E. L., Pedersen, B. K., & Krogh-Madsen, R. (2017). Ramadan model of intermittent fasting for 28 d had no major effect on body composition, glucose metabolism, or cognitive functions in healthy lean men. *Nutrition*, 37, 92–103. <https://doi.org/10.1016/j.nut.2016.12.015>
- Heart rate variability: Standards of measurement, physiological interpretation and clinical use. Task force of the european society of cardiology and the north american society of pacing and electrophysiology. (1996). *Circulation*, 93(5), 1043–1065.
- Hedelin, R., Kenttä, G., Wiklund, U., Bjerle, P., & Henriksson-Larsén, K. (2000). Short-term overtraining: Effects on performance, circulatory responses, and heart rate variability. *Medicine & Science in Sports & Exercise*, 32(8), 1480–1484. <https://doi.org/10.1097/00005768-200008000-00017>
- Hickey, K. C., Quatman, C. E., Myer, G. D., Ford, K. R., Brosky, J. A., & Hewett, T. E. (2009). Methodological report: Dynamic field tests used in an NFL combine setting to identify lower-extremity functional asymmetries. *Journal of Strength and Conditioning Research*, 23(9), 2500–2506. <https://doi.org/10.1519/JSC.0b013e3181b1f77b>
- Innes, E. (1999). Handgrip strength testing: A review of the literature. *Australian Occupational Therapy Journal*, 46(3), 120–140. <https://doi.org/https://doi.org/10.1046/j.1440-1630.1999.00182.x>
- Jahromi, S. R., Sahraian, M. A., Ashtari, F., Ayromlou, H., Etemadifar, M., Ghaffarpour, M., Mohammadianinejad, E., Nafissi, S., Nickseresht, A., Shaygannejad, V., Togha, M., Torabi, H. R., & Ziaie, S. (2014). Islamic fasting and multiple sclerosis. *BMC Neurology*, 14, 56–56. <https://doi.org/10.1186/1471-2377-14-56>
- James, D. V., Munson, S. C., Maldonado-Martin, S., & De Ste Croix, M. B. (2012). Heart rate variability: Effect of exercise intensity on postexercise response. *Research Quarterly for Exercise and Sport*, 83(4), 533–539. <https://doi.org/10.1080/02701367.2012.10599142>
- Kleiger, R. E., Miller, J. P., Bigger, J. T., Jr., & Moss, A. J. (1987). Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *American Journal of Cardiology*, 59(4), 256–262. [https://doi.org/10.1016/0002-9149\(87\)90795-8](https://doi.org/10.1016/0002-9149(87)90795-8)
- Kordi, R., Abdollahi, M., Memari, A. H., & Najafabadi, M. G. (2011). Investigating two different training time frames during ramadan fasting. *Asian Journal of Sports Medicine*, 2(3), 205–210. <https://doi.org/10.5812/asjsm.34774>
- Krustrup, P., Aagaard, P., Nybo, L., Petersen, J., Mohr, M., & Bangsbo, J. (2010). Recreational football as a health promoting activity: A topical review. *Scandinavian Journal of Medicine & Science in Sports*, 20(Suppl. 1), 1–13. <https://doi.org/10.1111/j.1600-0838.2010.01108.x>
- Krustrup, P., Randers, M. B., Andersen, L. J., Jackman, S. R., Bangsbo, J., & Hansen, P. R. (2013). Soccer improves fitness and attenuates cardiovascular risk factors in hypertensive men. *Medicine & Science in Sports & Exercise*, 45(3), 553–560. <https://doi.org/10.1249/MSS.0b013e3182777051>
- Kubota, Y., Chen, L. Y., Whitsel, E. A., & Folsom, A. R. (2017). Heart rate variability and lifetime risk of cardiovascular disease: The Atherosclerosis Risk in Communities Study. *Annals of Epidemiology*, 27(10), 619.e612–625.e612. <https://doi.org/10.1016/j.annepidem.2017.08.024>
- Lessan, N., & Ali, T. (2019). Energy metabolism and intermittent fasting: The Ramadan perspective. *Nutrients*, 11(5), 1192. <https://doi.org/10.3390/nut11051192>
- Lima-Silva, A. E., Bertuzzi, R. C. M., Pires, F. O., Fronchetti, L., Gevaerd, M. S., & De-Oliveira, F. R. (2010). A low carbohydrate diet affects autonomic modulation during heavy but not moderate exercise. *European Journal of Applied Physiology*, 108(6), 1133–1140. <https://doi.org/10.1007/s00421-009-1329-6>
- Luo, H., Newton, R. U., Ma'ayah, F., Galvão, D. A., & Taaffe, D. R. (2018). Recreational soccer as sport medicine for middle-aged and older adults: A systematic review. *BMJ Open Sport & Exercise Medicine*, 4(1), e000336. <https://doi.org/10.1136/bmjsem-2017-000336>
- MacDermid, J., Solomon, G., & Valdes, K., & American Society of Hand Therapists. (2015). *Clinical assessment recommendations*. American Society of Hand Therapists.
- Madsen, M., Krustrup, P., & Larsen, M. N. (2021). Exercise intensity during walking football for men and women aged 60+ in comparison to traditional small-sided football—a pilot study. *Managing Sport and Leisure*, 26(4), 259–267. <https://doi.org/10.1080/23750472.2020.1762508>

- Maughan, R. J., Fallah, J., & Coyle, E. F. (2010). The effects of fasting on metabolism and performance. *British Journal of Sports Medicine*, 44(7), 490–494. <https://doi.org/10.1136/bjsm.2010.072181>
- Mazurak, N., Gunther, A., Grau, F. S., Muth, E. R., Pustovoyt, M., Bischoff, S. C., Zipfel, S., & Enck, P. (2013). Effects of a 48-h fast on heart rate variability and cortisol levels in healthy female subjects. *European Journal of Clinical Nutrition*, 67(4), 401–406. <https://doi.org/10.1038/ejcn.2013.32>
- Melanson, E. L., & Freedson, P. S. (2001). The effect of endurance training on resting heart rate variability in sedentary adult males. *European Journal of Applied Physiology*, 85(5), 442–449. <https://doi.org/10.1007/s004210100479>
- Nachvak, S. M., Pasdar, Y., Pirsahab, S., Darbandi, M., Niazi, P., Mostafai, R., & Speakman, J. R. (2019). Effects of Ramadan on food intake, glucose homeostasis, lipid profiles and body composition composition. *European Journal of Clinical Nutrition*, 73(4), 594–600. <https://doi.org/10.1038/s41430-018-0189-8>
- Nematy, M., Alinezhad-Namaghi, M., Rashed, M. M., Mozhdehifard, M., Sajjadi, S. S., Akhlaghi, S., Sabery, M., Mohajeri, S. A., Shalaey, N., Moohebaty, M., & Norouzy, A. (2012). Effects of Ramadan fasting on cardiovascular risk factors: A prospective observational study. *Nutrition Journal*, 11, 69. <https://doi.org/10.1186/1475-2891-11-69>
- Nugraha, B., Ghashang, S. K., Hamdan, I., & Gutenbrunner, C. (2017). Effect of Ramadan fasting on fatigue, mood, sleepiness, and health-related quality of life of healthy young men in summer time in Germany: A prospective controlled study. *Appetite*, 111, 38–45. <https://doi.org/10.1016/j.appet.2016.12.030>
- Odemuyiwa, O., Farrell, T., Malik, M., Bashir, Y., Poloniecki, J., Ward, D. E., & Camm, A. J. (1991). The effect of age on the electrophysiological and autonomic correlates of sudden death after acute myocardial infarction. *Pacing and Clinical Electrophysiology*, 14(11 Pt 2), 2049–2055. <https://doi.org/10.1111/j.1540-8159.1991.tb02813.x>
- Persynaki, A., Karras, S., & Pichard, C. (2017). Unraveling the metabolic health benefits of fasting related to religious beliefs: A narrative review. *Nutrition*, 35, 14–20. <https://doi.org/10.1016/j.nut.2016.10.005>
- Prinsloo, G. E., Rauch, H. G. L., & Derman, W. E. (2014). A brief review and clinical application of heart rate variability biofeedback in sports, exercise, and rehabilitation medicine. *Physician and Sportsmedicine*, 42(2), 88–99. <https://doi.org/10.3810/psm.2014.05.2061>
- Ramadan, J. (2002). Does fasting during Ramadan alter body composition, blood constituents and physical performance? *Medical Principles and Practice*, 11(Suppl. 2), 41–46. <https://doi.org/10.1159/000066413>
- Ramadan, J. M., & Barac-Nieto, M. (2000). Cardio-respiratory responses to moderately heavy aerobic exercise during the Ramadan fasts. *Saudi Medical Journal*, 21(3), 238–244.
- Reddy, P., Dias, I., Holland, C., Campbell, N., Nagar, I., Connolly, L., Krstrup, P., & Hubball, H. (2017). Walking football as sustainable exercise for older adults—A pilot investigation. *European Journal of Sport Science*, 17(5), 638–645. <https://doi.org/10.1080/17461391.2017.1298671>
- Rodrigues, J. A. L., Yamane, A. C., Gonçalves, T. C. P., Kalva-Filho, C., Papoti, M., & Júnior, C. R. B. (2019). Fed and fasted states on heart rate variability, hemodynamic heart rate and blood pressure in adults submitted to moderate aerobic exercise. *International Journal of Cardiology & Heart & Vasculture*, 23, 100378. <https://doi.org/10.1016/j.ijcha.2019.100378>
- Sandrone, G., Mortara, A., Torzillo, D., La Rovere, M. T., Malliani, A., & Lombardi, F. (1994). Effects of beta blockers (atenolol or metoprolol) on heart rate variability after acute myocardial infarction. *American Journal of Cardiology*, 74(4), 340–345. [https://doi.org/10.1016/0002-9149\(94\)90400-6](https://doi.org/10.1016/0002-9149(94)90400-6)
- Sassi, R. H., Dardouri, W., Yahmed, M. H., Gmada, N., Mahfoudhi, M. E., & Gharbi, Z. (2009). Relative and absolute reliability of a modified agility T-test and its relationship with vertical jump and straight sprint. *Journal of Strength and Conditioning Research*, 23(6), 1644–1651. <https://doi.org/10.1519/JSC.0b013e3181b425d2>
- Scanlan, A. T., Wen, N., Pyne, D. B., Stojanovic, E., Milanovic, Z., Conte, D., Vaquera, A., & Dalbo, V. J. (2019). Power-related determinants of modified agility t-test performance in male adolescent basketball players. *Journal of Strength and Conditioning Research*, 35, 2248–2254. <https://doi.org/10.1519/jsc.0000000000003131>
- Schmidt, J. F., Hansen, P. R., Andersen, T. R., Andersen, L. J., Hornstrup, T., Krstrup, P., & Bangsbo, J. (2014). Cardiovascular adaptations to 4 and 12 months of football or strength training in 65- to 75-year-old untrained men. *Scandinavian Journal of Medicine & Science in Sports*, 24(Suppl. 1), 86–97. <https://doi.org/10.1111/sms.12217>
- Sezen, Y., Altıparmak, I. H., Erkus, M. E., Kocarslan, A., Kaya, Z., Gunebakmaz, O., & Demirbag, R. (2016). Effects of Ramadan fasting on body composition and arterial stiffness. *Journal of Pakistan Medical Association*, 66(12), 1522–1527.
- Shaffer, F., Meehan, Z. M., & Zerr, C. L. (2020). A critical review of ultra-short-term heart rate variability norms research. *Frontiers in Neuroscience*, 14, 594880. <https://doi.org/10.3389/fnins.2020.594880>
- Stein, P. K., Ehsani, A. A., Domitrovich, P. P., Kleiger, R. E., & Rottman, J. N. (1999). Effect of exercise training on heart rate variability in healthy older adults. *American Heart Journal*, 138(3), 567–576.
- Sweileh, N., Schnitzler, A., Hunter, G. R., & Davis, B. (1992). Body composition and energy metabolism in resting and exercising muslims during Ramadan fast. *Journal of Sports Medicine and Physical Fitness*, 32(2), 156–163.
- Temizhan, A., Dönderici, O., Ouz, D., & Demirbas, B. (1999). Is there any effect of Ramadan fasting on acute coronary heart disease events? *International Journal of Cardiology*, 70(2), 149–153. [https://doi.org/10.1016/s0167-5273\(99\)00082-0](https://doi.org/10.1016/s0167-5273(99)00082-0)
- Tornberg, J., Ikäheimo, T. M., Kiviniemi, A., Pyky, R., Hautala, A., Mäntysaari, M., Jämsä, T., & Korpelainen, R. (2019). Physical activity is associated with cardiac autonomic

- function in adolescent men. *PLOS ONE*, 14(9), e0222121. <https://doi.org/10.1371/journal.pone.0222121>
- Tsuji, H., Larson, M. G., Venditti, F. J., Jr, Manders, E. S., Evans, J. C., Feldman, C. L., & Levy, D. (1996). Impact of reduced heart rate variability on risk for cardiac events. The framingham heart study. *Circulation*, 94(11), 2850–2855. <https://doi.org/10.1161/01.cir.94.11.2850>
- White, D. W., & Raven, P. B. (2014). Autonomic neural control of heart rate during dynamic exercise: Revisited. *Journal of Physiology*, 592(12), 2491–2500. <https://doi.org/10.1113/jphysiol.2014.271858>
- Yusuf, S., Peto, R., Lewis, J., Collins, R., & Sleight, P. (1985). Beta blockade during and after myocardial infarction: An overview of the randomized trials. *Progress in Cardiovascular Diseases*, 27(5), 335–371. [https://doi.org/10.1016/s0033-0620\(85\)80003-7](https://doi.org/10.1016/s0033-0620(85)80003-7)
- Zainudin, S. B., Salle, D. D. A., & Aziz, A. R. (2021). Walking football during ramadan fasting for cardiometabolic and psychological health benefits to the physically challenged and aged populations. *Frontiers in Nutrition*, 8, 779863. <https://doi.org/10.3389/fnut.2021.779863>
- Zerguini, Y., Kirkendall, D., Junge, A., & Dvorak, J. (2007). Impact of Ramadan on physical performance in professional soccer players. *British Journal of Sports Medicine*, 41(6), 398–400. <https://doi.org/10.1136/bjsm.2006.032037>
- Zouhal, H., Saeidi, A., Salhi, A., Li, H., Essop, M. F., Laher, I., Rhibi, F., Amani-Shalamzari, S., & Ben Abderrahman, A. (2020). Exercise training and fasting: Current insights. *Open Access Journal of Sports Medicine*, 11, 1–28. <https://doi.org/10.2147/OAJSM.S224919>