



Isolated intraocular relapses of primary cerebral lymphomas: An loc network study

Nadia Younan, Carole Soussain, Sylvain Choquet, Nathalie Cassoux, Valérie Toutilou, Anna Schmitt, Olivier Chinot, Lucie Oberic, Gandhi Damaj, Roch Houot, et al.

► To cite this version:

Nadia Younan, Carole Soussain, Sylvain Choquet, Nathalie Cassoux, Valérie Toutilou, et al.. Isolated intraocular relapses of primary cerebral lymphomas: An loc network study. Hematological Oncology, 2022, 40 (5), pp.976-986. 10.1002/hon.3044 . hal-03716518

HAL Id: hal-03716518

<https://u-picardie.hal.science/hal-03716518v1>

Submitted on 16 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Isolated intraocular relapses of primary cerebral lymphomas: an LOC network study

Nadia Younan¹, Carole Soussain², Sylvain Choquet³, Nathalie Cassoux⁴, Valérie Touitou⁵, Anna Schmitt⁶, Olivier Chinot⁷, Lucie Oberic⁸, Gandhi Damaj⁹, Roch Houot¹⁰, Hervé Ghesquière¹¹, Kamel Laribi¹², Guido Ahle¹³, Luc Taillandier¹⁴, Jérôme Paillassa¹⁵, Emmanuel Gyan¹⁶, Fabrice Jardin¹⁷, Vincent Delwail¹⁸, Jean-Pierre Marolleau¹⁹, Adrian Tempescul²⁰, Philippe Agapé²¹, Marie Bourniquel²², Fabienne Vacheret²³, Ibrahim Jdid²⁴, Magali Le Garff-Tavernier²⁵, Denis Malaise²⁶, Agusti Alentorn¹, Khê Hoang Xuan¹, Caroline Houillier¹.

¹Neurology, Assistance Publique – Hôpitaux de Paris (APHP), Sorbonne Université, IHU, ICM, Groupe Hospitalier Pitié-Salpêtrière, Paris, France

²Hematology, Institut Curie, Site Saint-Cloud, France ; INSERM U932, Institut Curie, PSL Research University, Paris 75005, France

³Hematology, APHP, Groupe Hospitalier Pitié-Salpêtrière, Sorbonne Université, Paris, France

⁴Ophthalmology, Institut Curie, Site Paris, Université Paris V Descartes et PSL (Paris Science et Lettre), Paris, France

⁵Ophthalmology, APHP, Groupe Hospitalier Pitié-Salpêtrière, Sorbonne Université, Paris, France

⁶Hematology, Institut Bergonié, Bordeaux, France

⁷Neuro-oncology, Assistance Publique – Hôpitaux de Marseille (AP-HM), Timone, Marseille, France

⁸Hematology, Institut Universitaire du Cancer de Toulouse Oncopôle, Toulouse, France

⁹Hematology, Centre Hospitalier Universitaire de Caen, Université de Caen-Normandie, France

¹⁰Hematology, Centre Hospitalier Universitaire de Rennes, Université de Rennes, INSERM U1236, Rennes, France

¹¹Hematology, Centre Hospitalier Lyon Sud, Université Claude Bernard Lyon 1, Pierre-Bénite, France

¹²Hematology, Centre Hospitalier du Mans, Le Mans, France

¹³Neurology, Hôpital Pasteur – Hôpitaux civils de Colmar, France

¹⁴ Neurology, Centre Hospitalier Universitaire de Nancy, Nancy, France

¹⁵ Hematology, Centre Hospitalier Universitaire d'Angers, Angers, France

¹⁶ Hematology and cell therapy department, Centre Hospitalier Universitaire de Tours, Centre d'Investigations Cliniques INSERM U1517, Tours, France

¹⁷ Hematology, Centre Henri Becquerel, Rouen, France

¹⁸ Hematology, CHU de Poitiers, INSERM, CIC 1402, Centre d'Investigation Clinique, Université de Poitiers, Poitiers, France

¹⁹ Hematology, Centre Hospitalier Universitaire de Amiens, Amiens, France

²⁰ Hematology, Centre Hospitalier Universitaire de Brest, Brest, France

²¹ Oncology, Institut de Cancérologie de l'Ouest, Saint Herblain, France

²² Ophthalmology, Hôpital d'Instruction des Armées Sainte Anne, Toulon, France

²³ Hematology, Centre Hospitalier de Perpignan, Perpignan, France

²⁴ Hematology, Centre Hospitalier Régional d'Orléans, Orléans, France

²⁵ Service d'hématologie biologique, APHP, Groupe Hospitalier Pitié-Salpêtrière, Sorbonne Université, Paris, France

²⁶ Ophthalmology, Institut Curie, Paris ; PSL University, Inserm U1288, Laboratoire d'Imagerie Translationnelle en Oncologie, Orsay, France

Corresponding author: Caroline Houillier, MD, caroline.houillier@aphp.fr. Neuro-Oncology Department, Pitie Salpetriere Hospital, 47-83 boulevard de l'Hopital, 75013, Paris, France.
Phone : +33142164160, Fax : +33142160459

Data availability

All the data are available in the national French database of the “Lymphome Oculo-Cérébral” (LOC) network.

Funding information

The authors received no specific funding for this work.

Conflict of interest disclosure

The authors report no disclosures relevant to the manuscript.

Ethics approval statement

The study was approved by the Institutional Ethical Committee of the coordinating center on the 24/04/2018 and by the French “Commission Nationale de l’Informatique et des Libertés” (CNIL) (n°913170). This study was conducted in accordance with the Declaration of Helsinki.

Patient consent statement

All patients gave informed consent for the submission of their data to the database and their use.

Short running title: Isolated intraocular relapses of primary cerebral lymphomas

Keywords: primary CNS lymphoma, ocular lymphoma, relapse, prognosis

Abstract word count: 249 words

Main text word count: 2723 words

5 tables

1 figure

Supplementary data : 1 figure

Abstract

Most relapses of primary central nervous system lymphoma (PCNSL) occur in the brain and are associated with a poor prognosis. Isolated intraocular relapses (IIRs) are rare and poorly described.

We retrospectively selected from the French LOC database PCNSL patients who initially presented with cerebral localization and who experienced IIR during the course of the disease.

Of the 1472 patients included in the database, 55 patients presented an IIR. Their median age was 68 years, and median Karnofsky Performance Status 80. IL-10 levels in the aqueous humor and/or in the vitreous were increased in 42/46 patients. 45/55 patients received systemic chemotherapy, and 11/55 received high-dose chemotherapy with autologous stem cell transplantation (HCT-ASCT) as consolidation treatment. After a median follow-up of 69 months, 42/55 patients had relapsed, including 90% of the patients who did not receive HCT-ASCT at IIR and 40% of the patients who received HCT-ASCT at IIR ($p < 0.001$). The first relapse after the initial IIR was exclusively in the eye in 23/42 patients, and 29/42 patients had a subsequent brain relapse during the course of the disease. The median progression-free survival, brain-free survival and overall survival from IIR were 12.2, 48.6 and 57.1 months, respectively.

IIR is not exceptional in the course of PCNSL and deserves systematic ophthalmological follow-up. Its prognosis is much better than the prognosis of brain relapse, with an evolution close to that of primary vitreoretinal lymphoma. With the exception of patients who received HCT-ASCT at IIR, almost all patients subsequently relapsed, often with other IIRs.

Introduction

Primary central nervous system lymphoma (PCNSL) is a rare extranodal lymphoma affecting mostly elderly immunocompetent patients. It is located in the brain in more than 90% of cases. Intraocular involvement is a classic feature of the disease. At initial diagnosis, it can be either isolated, in rare cases¹, or associated with brain involvement in 10 to 20% of cases²⁻⁴. The association of an ocular location with a brain location does not seem to have an adverse prognostic impact³⁻⁵. However, it can be hard to eradicate and be a source of relapse.

The prognosis of PCNSL remains dismal, mainly due to a high risk of relapse during or after treatment. Due to unclear reasons, the disease remains confined to the central nervous system in most cases at relapse, with most of the relapses occurring in the brain⁶. Relapses in the brain can be associated with lymphomatous locations in the meninges, eye or spinal cord. Brain relapses have an adverse prognosis, with a median overall survival inferior to 12 months in most studies⁶⁻¹⁰, except for patients eligible for high-dose chemotherapy with autologous stem cell transplantation (HCT-ASCT)¹¹ or for late relapses when high-dose methotrexate (HD-MTX) can be rechallenged¹²⁻¹³.

Rarely, isolated intraocular relapses (IIRs) of primary cerebral lymphoma (PCL), without any brain involvement, can occur⁶. There are very limited data in the literature, and to our knowledge, no study specifically dedicated to this topic. The aim of the present study was to analyze, from a large cohort of PCNSL, patients with IIR at any time during their disease. The objectives were to search for risk factors for IIR and to describe cases of IIR, their management and their outcomes.

Methods

This work is based on an analysis of the French LOC network database, a nationwide database registering all newly diagnosed PCNSL since 2011 from 32 different centers in France, representing the main centers involved in PCNSL management. The database was approved by the Institutional Ethical Committee of the coordinating center and by the French “Commission Nationale de l’Informatique et des Libertés” (CNIL). All patients gave their informed consent to participate in the database and for the use of their data. This study was conducted in accordance with the Declaration of Helsinki. No private funds were used.

We selected adult immunocompetent patients who had a histologically proven diffuse large B-cell PCNSL with at least one cerebral localization at diagnosis and at least one isolated intraocular relapse (IIOR) at any point in the history of the disease. At the time of IIOR, cerebral involvement had to be excluded by cerebral MRI. Patients with asymptomatic meningeal involvement were not excluded. The patients were selected in July 2018, and data were analyzed in June 2021. The main characteristics and outcomes of these patients were compared to those of the PCNSL patients in the LOC network database with cerebral or oculocerebral brain relapse⁴.

Response to therapy was assessed according to the IPCG criteria¹⁴. IL-10 at diagnosis of IIOR was considered elevated with a cutoff of 30 pg/ml in the aqueous humor (AH) and 65 pg/ml in the vitreous according to previous publications in primary vitreoretinal lymphoma (PVRL)¹⁵⁻¹⁷. IL-10 in the AH was also monitored at the end of the treatment and during the follow-up at the discretion of the physicians. IL-10 levels were classified as detectable (≥ 2.5 pg/ml) or undetectable (< 2.5 pg/ml) for the follow-up. The chi-square test was used to test the associations between variables. The main outcome endpoints were overall survival (OS), progression-free survival (PFS) and brain-free survival (BFS). All of the endpoints were

calculated from the date of diagnosis of IIOR. PFS was defined as the time without relapse (whatever its location) or without death (whatever its cause). BFS was defined as the time without brain relapse. Survival rates were calculated using the Kaplan–Meier method. The log-rank test was used to test for equality of the PFS, BFS and OS distributions. A multivariate analysis, including the variables with significant prognostic value in univariate analysis, was performed with the multivariate Cox proportional hazards regression model. Two-sided p values <0.05 were considered significant. The statistical analyses were carried out with SPSS 17.0 software.

Results

Patients' characteristics at initial diagnosis

Of the 1472 PCNSL patients included in the LOC database at the time of patient selection, 615 relapsed, including 55 patients with IIOR from 23 centers who met the inclusion criteria (4% of the patients, 9% of the relapses). Their main characteristics are summarized in Table 1. At the time of the initial diagnosis, 49% of the patients had an associated ocular location. Of note, a search for an ocular location was not conducted at the time of the initial diagnosis for 11/55 (20%) of the patients. None of these patients had visual complaints. In first-line treatment, all patients received HD-MTX-based induction chemotherapy. None of them was treated with HCT-ASCT consolidation. Eighty percent of patients were in complete response (CR) at the end of first-line treatment.

Patients' characteristics at IIOR

The main characteristics at the time of diagnosis of IIOR are reported in Table 2. The median age at IIOR was 68 years (range 34-89), and the median KPS was 80 (range 50-100). IIOR occurred after a median duration of 16 months after the initial diagnosis. It was the first relapse in 48/55 patients and occurred during therapy in 5/48 patients. IIOR was asymptomatic in 5% of patients and revealed by visual acuity impairment in 86% (44/51). Ocular involvement was bilateral in 25/51 cases (49%). Hyalitis was observed in 44/48 (92%) of the cases. IL-10 levels in the AH and/or in the vitreous were increased in 42/46 patients (91%).

Treatments of IIOR

The treatments of the IIOR are summarized in Table 3. A total of 45/55 (82%) patients received systemic chemotherapy, including ten patients who received a combination with local

treatment. The treatment was based on ifosfamide in 13/55 cases, HD-MTX in 10/55 cases and temozolomide in 10/55 cases. Subsequently, 11/55 (20%) patients received thiotepa-based HCT-ASCT as consolidation treatment.

At the end of the treatment of the IIR, 27/54 (50%) were in CR. IL-10 was undetectable in the AH in 100% of the patients who were in CR/unconfirmed complete response (uCR) and detectable in 100% of the patients in stable disease (SD) or progressive disease (PD). Among the patients who received HCT-ASCT, 9/11 achieved CR, 1/11 progressed, and 1/11 died from toxic complications.

Outcome after IIR (Figure 1)

The median follow-up after IIR was 69 months (95% CI: 63-74), and 42/55 (76%) patients had a subsequent relapse(s). Their characteristics are summarized in Table 3. The first relapse after the initial IIR was exclusively in the eye in 23/42 patients (55%). Twenty-nine/42 (69%) patients had a subsequent brain relapse during the course of the disease after IIR diagnosis. Only 4/42 (10%) patients who did not receive HCT-ASCT after IIR diagnosis had no subsequent relapse, while 6/10 (60%) patients who received HCT-ASCT after IIR diagnosis never relapsed subsequently ($p < 0.001$). A total of 16/42, 17/42 and 10/42 patients received ibrutinib, lenalidomide or temozolomide, respectively, during subsequent relapse(s). The median PFS and BFS from the diagnosis of IIR were 12.2 months (95% CI: 8.7-15.7) and 48.6 months (27.3-63.9), respectively.

Twenty-six/55 (47%) patients died during the follow-up, due to cerebral progression of the PCNSL in 23 cases. The median OS from the diagnosis of IIR was 57.1 months (95% CI NR-NR), with 1-year, 2-year and 5-year OS rates of 87.3%, 78.1% and 49.9%, respectively.

Prognostic factors

The main prognostic factors for OS are indicated in Table 4. Age >70 years at IIR, KPS ≤70 at IIR, male sex and delay between initial diagnosis and IIR <18 months were associated with a shorter OS in univariate analysis. Only age >70 years and male sex remained significantly associated with a shorter OS in multivariate analysis.

There was no correlation between the levels of IL-10 in the vitreous or aqueous humor at IIR or the type of treatment at IIR (local vs. systemic vs. local+systemic) and PFS, BFS or OS. The use of HCT-ASCT at IIR was significantly associated with prolonged PFS ($p=0.005$) but was not associated with prolonged BFS or OS. However, there was a 6-year OS rate of 63% in patients treated with HCT-ASCT compared to 42% without HCT-ASCT (Supplemental Figure 1).

The median OS after IIR was 24 months when the first relapse after IIR was cerebral (+/- ocular), while it was 57.1 months when the first relapse after IIR was only ocular ($p=0.02$).

Comparison of patients in the LOC network database with cerebral or oculocerebral relapse

The cohort of IIR was compared to the patients with cerebral or oculocerebral relapse from the previously published cohort of the LOC network database⁴. The main results are indicated in Table 5. The median age was similar in both cohorts. The KPS was significantly higher at IIR than at first (ocular)cerebral relapse in the LOC database ($p<0.001$). The patients with IIR received significantly more active treatment ($p=0.002$) and more HCT-ASCT ($p=0.05$). The median PFS and median OS were significantly higher after IIR, in patients both older and younger than 70 years ($p<0.001$).

Discussion

To our knowledge, this work represents the first series dedicated to IIOR of primary cerebral lymphomas. Although rare³, this situation was not exceptional in our series, as it concerned 9% of the patients who relapsed during the course of their disease.

Even if ophthalmological involvement at diagnosis was far more frequent in IIOR than in classical PCNSL relapses, it is interesting to note that more than 50% of the patients had no history of ocular involvement at the initial diagnosis, either because no ophthalmological work-up was performed at that time (22% of the patients) or because this work-up was normal (30% of the patients). This underlines the importance of a systematic ophthalmological examination at the time of the diagnosis of a primary cerebral lymphoma, as recommended by the IPCG¹⁴, especially given that, generally, ocular symptoms are mild or even absent^{3,5}, whereas neurological symptoms are severe and are in the foreground. A systematic ophthalmological follow-up should also be recommended, at least in the first years after the end of the treatment. In the LOC network guidelines, an ocular assessment, including slit-lamp and fundus examination, is recommended once a year in PCNSL patients who had no ocular involvement at baseline. The site of disease in the brain at diagnosis was not recorded in this study but this analysis would be interesting to determine if some sites are associated with an increased risk of IIOR.

Of note, very few of the 55 patients with IIOR had received a consolidation treatment in first line : none received HCT-ASCT and 20% received WBRT. This reflects the practices in France at the time of the study : in PCNSL patients from the LOC network database diagnosed between 2011 and 2016, 6% and 15% received HCT-ASCT and WBRT in 1st line treatment, respectively⁴. Since the publication in 2017 and 2019 of the two main randomized trials addressing the question of WBRT vs HCT-ASCT as consolidation in 1st line

treatment^{15,16}, the use of HCT-ASCT in 1st line has increased a lot, so that there will surely be in the future some IIOR following HCT-ASCT. The eyes were not incorporated in the radiotherapy field of the patients who received WBRT in 1st line. We currently lack precise recommendations on the radiotherapy field in PCNSL, but these data support the incorporation of the eyes.

The IIOR clinical presentation was close to the PVRL clinical presentation, with mild visual acuity impairment and vitreous haze in most cases and frequent bilateral involvement^{1,3,5}. As in PVRL¹⁷⁻¹⁹, the IL-10 level in the vitreous or aqueous humor was a useful tool in the diagnosis and follow-up of IIOR, with 84% of patients with elevated IL-10 at the time of IIOR diagnosis, 100% of patients in CR/uCR after IIOR treatment with undetectable IL-10 and 100% of patients in SD or PD after IIOR treatment with elevated IL-10. As the ophthalmological evaluation according to the IPCG criteria is difficult and somewhat subjective, we recommend measuring IL-10 in the aqueous humor at least at baseline and at the end of the treatment of IIOR. A persistently increased IL-10 level despite an improved ophthalmological examination must result in close monitoring given the high risk of an early relapse¹. The search for a MYD88 mutation in the vitreous or in the aqueous humor, which is positive in 50 to 87% of cases²⁰⁻²², has become an important diagnostic tool in PVRL in the past years. Unfortunately, it was not available in the present study. Given the limited data on CSF IL-10 and the absence of data on CSF MYD88 mutation in the present study, we were not able to evaluate the prognostic impact of IL-10 and MYD88 mutation in the CSF in the setting of an IIOR.

Approximately 3/4 of the patients had subsequent relapses after IIOR. Most of the first subsequent relapses (55%) were also IIOR, while 69% of the patients relapsed in the brain during the follow-up, with a prolonged median BFS of 48.6 months. As in PVRL, ocular

lymphomatous locations appeared difficult to eradicate, with the final risk of brain relapse frequently leading to death.

As treatments were very heterogeneous in this retrospective cohort, it is difficult to determine the best therapeutic option in this setting. Temozolomide, lenalidomide and ibrutinib are three drugs administered orally, with a good tolerance profile even in elderly patients, that have proven efficacy against vitreoretinal lymphomas^{7,8,23}. They were offered in 20/55 (36%), 21/55 (38%), and 19/55 (34%) of the patients in this cohort, respectively, during the course of the disease. However, their exact place in relation to older and more toxic drugs such as MTX or ifosfamide is still unclear. The use of HCT-ASCT consolidation is standard in relapsed PCNSL who haven't received HCT-ASCT in first line treatment and who are young and fit enough for this treatment¹¹. In this cohort, the use of HCT-ASCT was significantly associated with improved PFS but not with BFS or OS. However, the rate of long-term survivors was much higher when HCT-ASCT was used, and almost all of the patients who did not receive HCT-ASCT at IIOR subsequently relapsed, while only 40% of the patients who received HCT-ASCT relapsed. HCT-ASCT appears to be the best chance to definitely cure the disease.

When compared to cerebral or oculocerebral relapses of PCNSL from the LOC database⁴, IIOR not surprisingly led to a much less severe clinical impairment (median KPS 80 vs. 60). This probably explains why active treatments (rather than palliative care) and HCT-ASCT could be offered more frequently in IIOR, despite the age being very similar in both cohorts. The prognosis of IIOR also appeared to be much better than the prognosis of brain relapses, with a PFS of 12.2 months (vs. 3.7 months, $p<0.001$) and a median OS of 57.1 months (vs. 6.8 months, $p<0.001$). In both cohorts, the majority of the patients died from the consequences of cerebral relapse. Many features probably contribute to this better prognosis: later relapses, known to be associated with better prognosis⁶; better KPS; more vigorous treatments; and a majority of IIOR at subsequent relapses after the first IIOR. It is interesting

to note that this favorable prognosis was also found in elderly patients for whom the prognosis at relapse is usually disastrous.

Conclusion

IIR of PCNSL with initial cerebral involvement is rare but not exceptional. It mainly occurs in patients with ocular involvement at initial diagnosis, but can also occur in patients without any ocular history, supporting the need for systematic ophthalmological follow-up of patients with PCNSL. The course of the disease is often similar to the course of PVRL, with many subsequent relapses still confined to the eyes and a high risk of final cerebral relapse. The prognosis of IIR appears much better than the prognosis of cerebral relapses, even in elderly patients. The optimal treatment remains unclear, but the better functional status at IIR allows for different treatment options ranging from classical drugs such as HD-MTX to newer drugs such as ibrutinib or lenalidomide and HCT-ASCT consolidation, which appears to be the best option to prevent subsequent relapses.

Acknowledgments

We gratefully thank the patients and their families for their participation in this study.

We acknowledge the research technicians of the LOC network (Bachir Aidoui, Diane Genet, Hassen Douzane, Yah-se Abada), Josiane Valero, all of the members of the LOC network and the Institut National of Cancer (INCa).

References

1. Lam M, Touitou V, Choquet S, Cassoux N, Ghesquières H, Kodjikian L, et al. Intravenous high-dose methotrexate based systemic therapy in the treatment of isolated primary vitreoretinal lymphoma: An LOC network study. *Am J Hematol*. 2021;96(7):823-833
2. Ferreri AJM, Blay JY, Reni M, Pasini F, Gubkin A, Tirelli U, et al. Relevance of intraocular involvement in the management of primary central nervous system lymphomas. International Extranodal Lymphoma Study Group (IELSG). *Ann Oncol*. 2002;13:531-538
3. Hong JT, Chae JB, Lee JY, Kim JG, Yoon YH. Ocular involvement in patients with primary CNS lymphoma. *J Neurooncol*. 2011;102:139-145
4. Houillier C, Soussain C, Ghesquières H, Soubeyran P, Chinot O, Taillandier L, et al. Management and outcome of primary CNS lymphoma in the modern era: An LOC network study. *Neurology*. 2020 Mar 10;94(10):e1027-e1039
5. Grimm SA, McCannel CA, Omuro AMP, Ferreri AJM, Blay JY, Neuwelt EA, et al. Primary CNS lymphoma with intraocular involvement: International PCNSL Collaborative Group Report. *Neurology*. 2008;71:1355-1360
6. Langner-Lemerrier S, Houillier C, Soussain C, Ghesquières H, Chinot O, Taillandier L, et al. Primary CNS lymphoma at first relapse/progression: characteristics, management, and outcome of 256 patients from the French LOC network. *Neuro Oncol*. 2016;18:1297-1303
7. Soussain C, Choquet S, Blonski M, Leclercq D, Houillier C, Rezai K, et al. Ibrutinib monotherapy for relapse or refractory primary CNS lymphoma and primary vitreoretinal lymphoma: Final analysis of the phase II 'proof-of-concept' iLOC study by the Lymphoma study association (LYSA) and the French oculo-cerebral lymphoma (LOC) network. *Eur J Cancer*. 2019;117:121-130

8. Ghesquieres H, Chevrier M, Laadhari M, Chinot O, Choquet S, Moluçon-Chabrot C, et al. Lenalidomide in combination with intravenous rituximab (REVRI) in relapsed/refractory primary CNS lymphoma or primary intraocular lymphoma: a multicenter prospective 'proof of concept' phase II study of the French Oculo-Cerebral lymphoma (LOC) Network and the Lymphoma Study Association (LYSA). *Ann Oncol*. 2019;30:621-628
9. Nayak L, Abrey LE, Drappatz J, Gilbert MR, Reardon DA, Wen PY, et al. Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma*. 2013;54:58-61
10. Arellano-Rodrigo E, López-Guillermo A, Bessell EM, Nomdedeu B, Montserrat E, Ghaus F. Salvage treatment with etoposide (VP-16), ifosfamide and cytarabine (Ara-C) for patients with recurrent primary central nervous system lymphoma. *Eur J Haematol*. 2003;70:219-224
11. Soussain C, Choquet S, Fourme E, Delgadillo D, Bouabdallah K, Ghesquière H, et al. Intensive chemotherapy with thiotepa, busulfan and cyclophosphamide and hematopoietic stem cell rescue in relapsed or refractory primary central nervous system lymphoma and intraocular lymphoma: a retrospective study of 79 cases. *Haematologica*. 2012 ;97:1751-1756
12. Plotkin SR, Betensky RA, Hochberg FH, Grossman SA, Lesser GJ, Nabors LB, et al. Treatment of relapsed central nervous system lymphoma with high-dose methotrexate. *Clin Cancer Res*. 2004;10:5643-5646.
13. Pentsova E, Deangelis LM, Omuro A. Methotrexate re-challenge for recurrent primary central nervous system lymphoma. *J Neurooncol*. 2014;117:161-165.
14. Abrey LE, Batchelor TT, Ferreri AJM, Gospodarowicz M, Pulczynski EJ, Zucca E, et al. Report of an international workshop to standardize baseline evaluation and response criteria for primary CNS lymphoma. *J Clin Oncol Off J Am Soc Clin Oncol*. 2005;23:5034- 5043.
15. Ferreri AJM, Cwynarski K, Pulczynski E, Fox CP, Schorb E, La Rosée P, et al. Whole-brain radiotherapy or autologous stem-cell transplantation as consolidation strategies after

- high-dose methotrexate-based chemoimmunotherapy in patients with primary CNS lymphoma: results of the second randomisation of the International Extranodal Lymphoma Study Group-32 phase 2 trial. *Lancet Haematol* 2017;4:e510-e523.
16. Houillier C, Taillandier L, Dureau S, Lamy T, Laadhari M, Chinot O, et al. Radiotherapy or Autologous Stem-Cell Transplantation for Primary CNS Lymphoma in Patients 60 Years of Age and Younger: Results of the Intergroup ANOCEF-GOELAMS Randomized Phase II PRECIS Study. *J Clin Oncol* 2019;37:823-833
17. Merle-Béral H, Davi F, Cassoux N, Baudet S, Colin C, Gourdet T, et al. Biological diagnosis of primary intraocular lymphoma. *Br J Haematol*. 2004;124:469- 473.
18. Cassoux N, Giron A, Bodaghi B, Tran THC, Baudet S, Davy F, et al. IL-10 measurement in aqueous humor for screening patients with suspicion of primary intraocular lymphoma. *Invest Ophthalmol Vis Sci*. 2007;48:3253- 3259.
19. Pochat-Cotilloux C, Bienvenu J, Nguyen A-M, Ohanessian R, Ghesquière H, Sève P, et al. Use of a threshold of interleukin-10 and il-10/il-6 ratio in ocular samples for the screening of vitreoretinal lymphoma. *Retina Phila Pa*. 2018;38:773- 781.
20. Raja H, Salomão DR, Viswanatha DS, Pulido JS. PREVALENCE OF MYD88 L265P MUTATION IN HISTOLOGICALLY PROVEN, DIFFUSE LARGE B-CELL VITREORETINAL LYMPHOMA. *Retina* 2016;36:624-628.
21. Bonzheim I, Giese S, Deuter C, Süsskind D, Zierhut M, Waizel M, et al. High frequency of MYD88 mutations in vitreoretinal B-cell lymphoma: a valuable tool to improve diagnostic yield of vitreous aspirates. *Blood* 2015;126:76-79.
22. Belhouachi N, Xochelli A, Boudjoghra M, Lesty C, Cassoux N, Fardeau C, et al. Primary vitreoretinal lymphomas display a remarkably restricted immunoglobulin gene repertoire. *Blood Adv* 2020;4:1357-1366.

23. Baron M, Belin L, Cassoux N, Fardeau C, Blaizeau M, Soussain C, et al. Temozolomide is effective and well tolerated in patients with primary vitreoretinal lymphoma. *Blood*. 2020 ;135:1811-1815.

Table 1: Patient characteristics at initial diagnosis, details of first-line treatment

Table 2: Patient characteristics at IIOR

Table 3: Treatments of IIOR and outcomes

Table 4: Prognostic factors in terms of OS (uni- and multivariate analysis)

Table 5: Comparison between characteristics of IIOR and characteristics of first (oculo)cerebral relapses in the whole LOC network database cohort

Figure 1: Progression-free survival, brain-free survival and overall survival after IIOR

Supplemental Figure 1: Progression-free survival, brain-free survival and overall survival after IIOR according to the use of HCT-ASCT at IIOR or not

Table 1: Patient characteristics at initial diagnosis, details of first-line treatment

N	55
Age : median (range)	67 (33-88)
Sex ratio (M/W)	25/30
KPS: median (range)	70 (40-100)
Associated ocular disease	
Yes (%)	27/55 (49%)
No (%)	17/55 (31%)
Unknown (%)	11/55 (20%)
Cerebral lesions	
Unique (%)	23/48 (48%)
Multiple (%)	25/48 (52%)
Meningeal involvement (%)	5/30 (17%)
1 st line treatment	
HD MTX-based CT	55/55 (100%)
Rituximab	28/53 (53%)
Consolidation treatment	
HCT-ASCT	0/55 (0%)
WBRT	11/55 (20%)
With eyes incorporated in the RT field	0
With eyes not incorporated in the RT field	10
Unknown	1
Response to 1st line treatment (including WBRT)	
CR/uCR	44/55 (80%)
PR	4/55 (7%)
SD	0/55 (0%)
PD	7/55 (13%)
PD in the brain	2/55 (4%)
IIOR	5/55 (9%)

Table 2: Patient characteristics at IIOR

N	55
Median age (range)	68 (34-89)
Median KPS (range)	80 (50-100)
Median delay from initial diagnosis (months) (range)	16 (3-101)
In patients with ophthalmological involvement at baseline	13 (3-51)
In patients without ophthalmological involvement at baseline	18 (8-101)
Time of IIOR	
First relapse	48/55 (87%)
- refractory	5/48 (10%)
- relapsed	43/48 (90%)
Second relapse	7/55 (13%)
-refractory	2/7 (29%)
-relapsed	5/7 (71%)
Symptoms	
Visual acuity impairment (%)	44/51 (86%)
No symptom (%)	3/55 (5%)
Vitreous haze	44/48
Unilateral ocular involvement	26/51 (51%)
Bilateral ocular involvement	25/51 (49%)
Median best corrected visual acuity of affected eyes at diagnosis, logMAR (range)	+0.3 (0-1.3)
IL10 and IL6 dosage in AH and/or vitreous	
Number of data available	46/55 (84%)
Median IL10 in AH (pg/ml) (range)	214 (4-5000)
Median IL10 in vitreous (pg/ml) (range)	720 (0-10453)
Increased IL10 in AH and/or vitreous (%)	42/46 (91%)
IL10/IL6 >1 in AH and/or vitreous	33/36 (92%)
CSF analysis	
Lumbar puncture performed	35/52 (67%)
Lymphomatous cells (cytology or flow cytometry)	2/35 (6%)
No lymphomatous cells	33/35 (94%)
CSF IL-10	
Detectable	4/10 (40%)
Undetectable	6/10 (60%)

Table 3: Treatments of IIOR and outcomes

N	55
Type of treatment	
Only systemic CT	35/55 (64%)
Only local treatment	8/55 (15%)
IVT MTX	6
Ocular RT	2
Both systemic and local treatment	10/55 (18%)
Only steroids	1/55 (2%)
No treatment	1/55 (2%)
Systemic treatment	45/55 (82%)
ifosfamide-based	13/55 (24%)
methotrexate-based	10/55 (20%)
temozolomide alone	10/55 (20%)
lenalidomide	4/55 (7%)
AraC-based	3/55 (5%)
ibrutinib	3/55 (5%)
other	2/55 (4%)
rituximab	19/55 (35%)
Local treatment	18/55 (33%)
IVT MTX	13/55 (24%)
Ocular RT	5/55 (9%)
Thiotepa-based HCT-ASCT	11/55 (20%)
Median age at IIOR (range)	60 (50-73)
Median KPS at IIOR (range)	80 (70-90)
Response to treatment	
CR/uCR	27/54 (50%)
PR	0
SD	1/54 (2%)
PD	24/54 (44%)
PD only in the eye	15/54 (28%)
PD in the brain +/- in the eye	9/54 (17%)
Not evaluable	2/54 (4%)
IL-10 in the AH at the end of the treatment	N=33
In patients with CR/uCR in the eye	
Detectable	0
Undetectable	18/18 (100%)
In patients with SD/PD in the eye	
Detectable	15/15 (100%)
Undetectable	0
Number of patients who relapsed after the IIOR	42/55 (76%)
In patients with HCT-ASCT at IIOR	4
In patients with no HCT-ASCT at IIOR	38
Number of patients who never relapsed after IIOR	13/55 (24%)
Early death after IIOR	3
In patients with HCT-ASCT at IIOR	6

In patients with no HCT-ASCT at IIOR	4
Localization of the first relapse post-IIOR	
- Only brain	14/42 (33%)
- Only eye	23/42 (55%)
- Both brain and eye	5/42 (12%)
Localization of relapses throughout the disease post IIOR	
- Only Brain	12/42 (29%)
- Only eye	13/42 (31%)
- Both brain and eye	17/42 (40%)
HCT-ASCT at subsequent relapses after IIOR	4/42 (10%)
Death N (%)	26/55 (47%)
Cause of death	
Cerebral progression of the lymphoma	23
Treatment-related toxicity	2
Other	1

Table 4: Prognostic factors in terms of OS (uni- and multivariate analysis)

	N	Univariate analysis		Multivariate analysis	
		Median OS	p	HR	P
Age at IIR <70 years	32	NR	0.01	2.9	0.02
Age at IIR >70 years	23	37.1			
KPS at IIR ≤70	8	24	0.05	0.4	0.07
KPS at IIR >70	43	NR			
Female sex	30	NR	0.03	2.5	0.04
Male sex	25	45.9			
Delay diag-IIR<18 months	35	40.1	0.01	0.4	0.14
Delay diag-IIR > 18 months	20	NR			
IIR at 1 st relapse	48	57.1	0.3		
IIR at 2d relapse	7	NR			
Unilateral IIR	26	NR	0.2		
Bilateral IIR	25	55.6			
IL-10 in AH or vitreous< median value	23	57.1	0.9		
IL-10 in AH or vitreous≥ median value	22	51.1			
Type of treatment			0.3		
Systemic	35	NR			
Local	8	40.4			
Local + systemic	10	34.1			
Type of systemic treatment			0.1		
ifosfamide-based	13	37.5			
methotrexate-based	10	NR			
temozolomide alone	10	37.1			
Rituximab at IIR	19	NR	0.3		
No rituximab at IIR	36	51.1			
HCT-ASCT at IIR	11	NR	0.4		
No HCT-ASCT at IIR	44	55.6			
Localization of 1st relapse post IIR			0.02		
Brain+/- eye	19	24			
Only eye	23	57.1			


NR : not reached

Table 5: Comparison between characteristics of IIOR and characteristics of first (oculo)cerebral relapses in the whole LOC network database cohort


	IIOR	First relapses in the whole LOC network database cohort	p
N	55	471	
Median age (years)	68	69	NS
Sex ratio M/W	0.8	1.2	NS
OPH involvement at baseline			
Yes	27 (49%)	56 (13%)	<0.001
No	17 (31%)	233 (52%)	
No work-up	11 (20%)	157 (35%)	
1 st line of treatment			
HD-MTX based CT	56 (100%)	451 (96%)	NS
Median time between initial diagnosis and relapse (months)	16	8	
Refractory to previous treatment	7	232	<0.001
Not refractory to previous treatment	48	239	
Median KPS at relapse	80	60	<0.001
KPS ≥ 70	50	105	
KPS < 70	1	346	
Treatment at relapse			
Active treatment	54 (98%)	378 (82%)	0.002
Palliative care	1 (2%)	82 (18%)	
HCT-ASCT (at any time of the follow-up after relapse)	15 (27%)	76 (17%)	0.05
No HCT-ASCT	40 (73%)	384 (83%)	
Median PFS after relapse (months)	12.2	3.7	<0.001
Median OS after relapse (months)	57.1	6.8	<0.001
In patients > 70 years	37.1	4.6	<0.001
In patients < 70 years	NR	8.4	<0.001

NS : not significant

Progression-free survival



Brain-free survival



Overall survival

