



## Supplemental Online Content

<b>efigure 1. Symptom combinations among the 194 patients with three or more symptoms at M12 visit .....</b>	<b>2</b>
<b>Supplementary Table 1. Associations of presence of <math>\geq 3</math> symptoms at M12 with MRC scale, modified Rankin scale, HADS at M12 .....</b>	<b>2</b>
<b>Supplementary Table 2. Associations of the HADS Anxiety scores with age, sex, hospitalisation in intensive care unit, dyspnea, back to work .....</b>	<b>3</b>
<b>efigure 2. Observed proportions (and their 95% CI) of three or more persistent symptoms at M12 according to the combination of the risk factors found in the multivariate model in women and in men separately .....</b>	<b>4</b>


**figure 1. Symptom combinations among the 194 patients with three or more symptoms at M12 visit**



**In women with  $\geq 3$  symptoms at M12 (n=98)**



**In men with  $\geq 3$  symptoms at M12 (n=96)**



**Supplementary Table 1. Associations of presence of  $\geq 3$  symptoms at M12 with MRC scale, modified Rankin scale, HADS at M12**


	Missing	<3 symptoms at M12 N=516	$\geq 3$ symptoms at M12 N=194	Bivariate analysis	
				OR [95% CI]	p-value
MRC Scale <48 at M12 visit – n (%)	227	4 (1.2%)	4 (2.7%)	2.30 [0.54, 9.84]	0.24
Simplified MRS $\geq 2$ at M12 visit – n (%)	234	43 (13.0%)	60 (41.7%)	4.80 [3.04; 7.65]	<0.001
HADS – Anxiety $\geq 11$ – n (%)	291	22 (7.7%)	37 (27.4%)	4.50 [2.55; 8.11]	<0.001
HADS – Depression $\geq 11$ – n (%)	291	15 (5.3%)	20 (14.9%)	3.16 [1.57; 6.49]	0.001
SF12 – Impaired physical HRQL – n (%)	309	110 (40.6%)	85 (65.4%)	2.76 [1.80; 4.30]	<0.001
SF12 – Impaired mental HRQL – n (%)	309	68 (25.1%)	58 (44.6%)	2.40 [1.55; 3.75]	<0.001

**Supplementary Table 2. Associations of the HADS Anxiety scores with age, sex, hospitalisation in intensive care unit, dyspnea, back to work**


	Missing	M12 HADS –	M12 HADS –	Bivariate analysis	
		Anxiety <11 N=361	Anxiety ≥11 N=59	OR [95% CI]	p-value
Age, Median [IQR], y	0	59 [52 ; 69]	58 [51 ; 67]	0.99 [0.97; 1.02]	0.56
Female gender, n (%)	0	123 (34.1%)	33 (55.9%)	<b>2.46 [1.41; 4.32]</b>	<b>0.002</b>
Intensive care unit during acute phase, n (%)	64	96 (31.5%)	14 (27.5%)	0.82 [0.41; 1.56]	0.57
Not back to work at M12, n (%)	208	42 (22.7%)	12 (44.4%)	<b>2.72 [1.17; 6.27]</b>	<b>0.018</b>
Dyspnea at M12, n (%)	2	115 (31.9%)	36 (62.1%)	<b>3.49 [1.98; 6.27]</b>	<b>&lt;0.001</b>

**efigure 2. Observed proportions (and their 95% CI) of three or more persistent symptoms at M12 according to the combination of the risk factors found in the multivariate model in women and in men separately**

**WOMEN**



**MEN**



The proportions of ≥3 persistent symptoms at M12 are reported with their 95%CI. The number of patients with and without M12 visit in each of the combinations are reported.