

# Occupational exposures in the rubber tire industry and risk of cancer: a systematic review

Sylvain Chamot, Marion Delzard, Lea Leroy, Gwen Marhic, Olivier Ganry

## ► To cite this version:

Sylvain Chamot, Marion Delzard, Lea Leroy, Gwen Marhic, Olivier Ganry. Occupational exposures in the rubber tire industry and risk of cancer: a systematic review. International Archives of Occupational and Environmental Health, 2023, 10.1007/s00420-023-01972-x . hal-04083787

## HAL Id: hal-04083787 https://u-picardie.hal.science/hal-04083787

Submitted on 11 May 2023  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Occupational Exposures in the Rubber Tire Industry and Risk of Cancer: a Systematic Review

Authors: Sylvain Chamot<sup>1,2</sup>, Marion Delzard<sup>3</sup>, Lea Leroy<sup>1</sup>, Gwen Marhic<sup>1,2</sup>, Olivier Ganry<sup>4</sup>

Affiliations:

<sup>1</sup> Regional Center for Occupational and Environmental Pathologies of Hauts-de-France, Amiens University Hospital, Amiens, France

<sup>2</sup> Péritox (UMR\_I 01); UPJV/INERIS; UPJV, Amiens, France

<sup>3</sup> Department of General Medicine, Amiens University Hospital, Amiens, France.

<sup>4</sup> Cancer Registry of the Somme, Department of Epidemiology, Sanitation, and Public Health, Amiens University Hospital, Amiens, France

Corresponding author: Sylvain Chamot, <u>chamot.sylvain@chu-amiens.fr</u> (ORCID : 0000-0001-8769-8090)

Word count: 3920 words Abstract: 212 words Figures: 2

Tables: 1

Number of references: 32

#### Abstract:

**Objective:** To investigate the association between cancer and the rubber tire industry using indicators used to estimate exposure, such as duration or historical time period of exposure.

**Methods:** A systematic review using MEDLINE was carried out. The Preferred Reporting Items for Systematic Reviews (PRISMA) statement guided the systematic review. The Risk of Bias in Non-Randomized Studies of Exposures (ROBINS-E) was used to analyze biases in the selected studies.

**Results:** A total of 240 articles were initially selected. Of these, 17 were included in this systematic review of 15 different cohorts and one case-control study. Five cohorts were composed of rubber industry employees including some who who worked in the rubber tire industry, whereas the other 10 cohorts were composed exclusively of rubber tire industry employees. An increased risk of lung/respiratory cancer is possible in the oldest time periods without duration of exposure showing any association according to the analysis. The work areas of mixing and milling as well as compounding seem to be more specifically associated with this risk, which could be expected because of the exposure to fumes containing carcinogenic agents.

**Conclusions:** Having worked in the rubber tire industry in the past, especially in milling, mixing as well as compounding may be associated with an increased risk of lung/respiratory cancer.

**Keywords:** Rubber, Occupational Diseases, Neoplasms, Humans, Cohort Studies, Case-control studies

## 1 Introduction

Occupational exposures related to rubber industry activities are classified as Group 1 by the International Agency for Research on Cancer (IARC).

The first evaluation by IARC classifying the rubber industry as a carcinogen was in 1982, and it was confirmed in 1987 and 2012 (International Agency for Research on Cancer, 1982, 1987, 2012). There is sufficient evidence of carcinogenicity for lung, bladder, and stomach cancers and also for leukemia and lymphoma. There is limited evidence of carcinogenicity for esophageal, laryngeal, and prostate cancer.

The rubber industry uses hundreds of different chemicals. This wide variety has never allowed IARC to establish the precise role of each individual exposure in the development of cancer. We know that several compounds known to be carcinogens have been used in this industry. On the other hand, carcinogens were formed during the manufacturing processes. Additionally, tools and machines containing asbestos were used in these factories. Although there are common processes among the different branches of the rubber industry, there is some heterogeneity in terms of exposure. In the curing processes, for example, three main techniques are used in the plants: compression/injection, UHF microwaves, and nitrite salt baths. The salt baths expose workers much more to N-nitrosamines than the other techniques (Héry, 2009). Similarly, some plants in the rubber industry have workplaces dedicated to the production of synthetic rubber. Plants producing styrene-butadiene synthetic rubber use 1-3 butadiene as a raw material, sometimes with high exposure, whereas plants supplied with an external source of styrene-butadiene rubber have lower exposure to 1-3 butadiene (Weiland et al., 1998). The European project ExAsRUB highlights higher exposures to different agents for employees working in the general rubber goods (GRG) industry when compared with workers in the tire manufacturing industry (Hidajat et al., 2019). Rubber fume concentrations (Tire geometric mean: 0.17 mg/m<sup>3</sup>; GRG geometric mean: 0.39 mg/m<sup>3</sup>; p < 0.05) and the sum of n-nitrosamines (Tire geometric mean:  $0.24 \mu g/m^3$ ; GRG geometric mean:  $1.06 \mu g/m^3$ ; p < 0.05) were lower in the tire manufacturing industry than in the GRG industry. These data are consistent with an earlier British study (Dost et al., 2000) where rubber fume concentrations were lower in the tire industry than in the GRG industry (Tire mean: 0.23 mg/m; GRG mean: 0.53 mg/m<sup>3</sup>).

This heterogeneity in processes and exposures, especially between the GRG industry and the tire manufacturing industry, has led most authors who have conducted studies in this field to identify rubber tire manufacturing plants separately in order to refine the data and obtain a more homogeneous group in terms of exposure.

Alder et al. and Boniol et al. conducted meta-analyses in 2006 and 2017 on the excess risk of cancer in employees of the rubber industry, with an analysis of the tire manufacturing industry (Alder et al., 2006; Boniol et al., 2017). They did not find any significant increase in risk for different cancers in the tire manufacturing industry. However, neither of the articles addressed the more precise analyses carried out in the included studies, such as risk by work area or years of exposure. No analysis of risk of bias is presented in their work. It appeared, instead, that great disparities were present within the same company. For example, employees assigned to curing had much higher exposure to carcinogens contained in vulcanization fumes. Moreover, N-nitrosamine and aromatic amines exposure, due to technical improvements, dropped as the industry was modernized (Dost et al., 2000; Goutet, 2007).

In order to more fully elaborate on the topic of the meta-analyses of Alder and Boniol (Alder et al., 2006; Boniol et al., 2017) ,we carried out a systematic review on the risk of cancer in the rubber tire manufacturing industry with the aim of presenting the results by type of work area, years of exposure, duration of exposure, and other exposure information. We were interested in cancer site studies associated with sufficient evidence of carcinogenicity in the rubber industry according to IARC.

## 2 <u>Methods</u>

#### **Research questions**

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used to guide this systematic review (Page et al., 2021). This systematic review aimed to answer this question: Is there an increased risk of cancer for rubber tire industry employees if we examine the data characterizing their exposure?

#### Literature search strategy

Two reviewers searched MEDLINE (via PubMed) for all articles available until the last search date, July 1, 2022. The sequence of terms that was used to search on PubMed was built with the Yale MeSH Analyzer (Grossetta Nardini & Wang, 2022) and validated with five clearly eligible studies (De Vocht et al., 2009; Mirabelli et al., 2012; Pira et al., 2012; Wilczyńska et al., 2001; Wingren & Axelson, 2007) and was: ("Rubber"[Mesh]) AND ("Occupational Diseases"[Mesh] OR "Neoplasms"[Mesh]) AND ("Humans"[Mesh]) AND ("Cohort Studies"[Mesh] OR "Case-Control Studies"[Mesh] OR "Neoplasms/epidemiology"[Majr] OR "Neoplasms/mortality"[Majr]). Reference checking of included articles was done to identify additional studies that were missed. We also examined the reference lists of the two meta-analyses cited above (Alder et al., 2006; Boniol et al., 2017). A complementary search was performed on WEB OF SCIENCE with the algorithm "(tire or tyre) and cancer".

#### **Eligibility criteria**

We excluded all articles that were not on human epidemiological cohorts and case-control studies, that did not relate to the rubber tire industry (or in which results for this industry were not reported) as well as studies for which no analysis of statistical significance was performed. We included both incidence and mortality studies and did not specify a minimum duration of follow-up. We did not put any restriction on the year of publication, but they had to be articles published in a scientific journal in English.

#### **Selection process**

Two reviewers screened each record, and each report that was retrieved, by working together was shared. Agreement was then reached on whouch articles would be retained. The first

selection was made on the basis of the title and the abstract, and the second after reading the remaining full-text articles.

#### **Study selection**

The search algorithm identified 239 articles. We identified one additional article of interest from the previously cited meta-analyses that was not over-referenced on MEDLINE (Chen et al., 1997). Of these 240 articles, 210 were excluded for not meeting the inclusion criteria or for being identified as an older article from the same study (and were therefore a duplicate). Thus, 30 studies remained to be screened by full-text reading. Finally, 17 articles (Beall et al., 2007; Bernardinelli et al., 1987; Boniol et al., 2016; Chen et al., 1997; Coggon et al., 1990; De Vocht et al., 2009; McElvenny et al., 2018; McMichael et al., 1974; Mirabelli et al., 2012; Negri et al., 1989; Pira et al., 2012; Sorahan et al., 1989; Veys, 2004; Vlaanderen et al., 2013; Wang et al., 1984; Wilczyńska et al., 2001; Wingren & Axelson, 2007) were included in this systematic review, including two on the same cohort with quite different analyses (De Vocht et al., 2009; Wilczyńska et al., 2001) and only one case-control study (Coggon et al., 1990). Of the 13 articles not included, one did not perform a statistical analysis, one only focused on solvents, and in eleven it was not possible to separate tires from other rubber goods. The search on WEB OF SCIENCE allowed to add a complementary study among the 625 results found, bringing to 18 the number of articles studied (Rubin & Clapp, 2012). The flow chart of the systematic review is described in Fig 1 and data on retained and excluded articles are available as Supplementary data. We did not find any relevant case series.

For each study selected, we examined the results and extracted the data according to the approaches used to estimate the risk (duration of exposure, hiring period, etc.).

Of the eighteen articles, seven were not cited in the last meta-analysis (Boniol et al., 2017). Three were not included (De Vocht et al., 2009; Mirabelli et al., 2012; Wilczyńska et al., 2001) because they overlapped with another article (Boniol et al., 2016), one was an update of a previous study published after the release of the meta-analysis (Fox & Collier, 1976; McElvenny et al., 2018), and three did not appear in the search results of Boniol et al. (Beall et al., 2007; Chen et al., 1997; Rubin & Clapp, 2012).

#### Assessment of risk of bias in individual studies

A reviewer assessed the risk of bias in the selected studies. The Risk of Bias in Non-Randomized Studies of Exposures (ROBINS-E) tool was used for the risk of bias assessment (ROBINS-E Development Group, 2022). The seven domains of ROBINS-E were: 1) bias due to confounding, 2) bias in measurement of the exposure, 3) bias in selection of participants into the study, 4) bias due to post-exposure interventions, 5) bias due to missing data, 6) bias in measurement of outcomes, and 7) bias in the selection of the reported results. The 4th domain was not relevant to the studies in the review and was not considered. The judgment for each ROBINS-E item was defined as "Low Risk of Bias", "Some Concerns", "High Risk of Bias", or "Very High Risk of Bias". For the overall Risk-of-Bias Rating, another judgment was added: "Low Risk of Bias, except for concerns about residual confounding". In order to be consistent with other publications, we reformulated these judgments to read: "Low", "Moderate", "Serious", or "Critical". Risk of bias assessment data are available as Supplementary data.

## 3 <u>Results</u>

All studies were published from 1974 to 2018. The case-control study took place in England, with 15 cases in the rubber tire industry (Coggon et al., 1990). The cohort sample size varied from 3,425 (Beall et al., 2007) to 17,636 (De Vocht et al., 2009; Wilczyńska et al., 2001). Four studies were performed in Italy (Bernardinelli et al., 1987; Mirabelli et al., 2012; Negri et al., 1989; Pira et al., 2012), four in Great-Britain (Coggon et al., 1990; McElvenny et al., 2018; Sorahan et al., 1989; Veys, 2004), two in China (Chen et al., 1997; Wang et al., 1984), three in the USA (Beall et al., 2007; McMichael et al., 1974; Rubin & Clapp, 2012), one in Germany (Vlaanderen et al., 2013), one in Poland (De Vocht et al., 2009; Wilczyńska et al., 2001), and one in Sweden (Wingren & Axelson, 2007). One publication (Boniol et al., 2016) includes the results of several cohorts (De Vocht et al., 2009; Mirabelli et al., 2012; Wilczyńska et al., 2001) of employees first employed since 1975, and its results are from German, Italian, Polish, Swedish, and English cohorts.

#### Approaches used to estimate risk

Five studies were conducted in companies that were not all in the rubber tire sector (Boniol et al., 2016; Chen et al., 1997; McElvenny et al., 2018; Sorahan et al., 1989; Vlaanderen et al., 2013), whereas the other twelve were conducted in the rubber tire industry only (Beall et al., 2007; Bernardinelli et al., 1987; Coggon et al., 1990; De Vocht et al., 2009; McMichael et al., 1974; Mirabelli et al., 2012; Negri et al., 1989; Pira et al., 2012; Rubin & Clapp, 2012; Veys, 2004; Wang et al., 1984; Wilczyńska et al., 2001; Wingren & Axelson, 2007).

Among the twelve studies conducted exclusively in the rubber tire industry, seven did an analysis of risk by duration of employment (Beall et al., 2007; Coggon et al., 1990; Mirabelli et al., 2012; Negri et al., 1989; Pira et al., 2012; Wilczyńska et al., 2001; Wingren & Axelson, 2007), four by time elapsed since first employment (Beall et al., 2007; Coggon et al., 1990; Mirabelli et al., 2012; Pira et al., 2012), two by the end date of employment and age at the start of employment (Negri et al., 1989; Pira et al., 2012), three by time period at first employment (Negri et al., 1989; Pira et al., 2012), three by time period at first employment (Negri et al., 1989; Pira et al., 2012), and four by work area (Negri et al., 1989; Wang et al., 1984; Wilczyńska et al., 2001; Wingren & Axelson, 2007), and two by sex (Rubin & Clapp, 2012; Wilczyńska et al., 2001). One study used an exposure matrix for aromatic amines and inhalable aerosols (De Vocht et al., 2009).

The characteristics of the eligible studies in this systematic review are shown in Table 1.

#### **Risk of bias**

The summary of the risk of bias in the eight included studies is shown in Fig 2. Every study had a serious risk of bias due to confounding factors except one, which focused on digestive cancers (Chen et al., 1997), because none of them took into account the exposure to the major extraoccupational risk factor of smoking or other factors such as exposure to benzene or asbestos for example.

An Italian (Negri et al., 1989), a Chinese (Wang et al., 1984), and an Americain cohorts (Rubin & Clapp, 2012)and the case-control study (Coggon et al., 1990) had a critical risk of bias due to the selection of the reported results, classification of exposures, a great risk of bias, and a bias due to missing data, respectively. Threestudies (Boniol et al., 2016; Chen et al., 1997; De Vocht et al., 2009; Wilczyńska et al., 2001) presented a serious overall risk of bias, and six (Bernardinelli et al., 1987; McMichael et al., 1974; Mirabelli et al., 2012; Pira et al., 2012; Sorahan et al., 1989; Vlaanderen et al., 2013) a moderate risk of bias. In the remaining four studies, the risk of overall bias was low except for confounding factors.

#### Main findings

Table 1 shows the main findings of the studies, showing the results according to duration of employment, work area, and other results.

Among the twelve studies conducted exclusively in the rubber tire industry, no significant increase in risk was found to be associated with duration of employment, with some exceptions. For example, Wingren et al.'s. study of SIR for tracheal, bronchial, and lung cancer in employees exposed from 2.5 to 10 years and for those exposed for more than 10 years, there was an increase (SIR = 1.75 [1.04-2.77], and 1.78 [1.09-2.75, respectively) (Wingren & Axelson, 2007).

No association was found by studying the time elapsed since first employment (Beall et al., 2007; Mirabelli et al., 2012; Pira et al., 2012), the time since last employment, the age at when first employed, and the time period of first employment (Negri et al., 1989; Pira et al., 2012; Rubin & Clapp, 2012).

Four studies have examined the work area of employees.Wilczyńska 2001 et al. (Wilczyńska et al., 2001) did not find any association whereas Wingren et al. (Wingren & Axelson, 2007) found an association between respiratory cancer and compounding, mixing, milling, and maintenance (SIR for tracheal, bronchial, and lung cancers for employees in compounding and

mixing: 2.18 [1.31-3.4]; SIR for tracheal,, bronchial, and lung cancers for employees in milling: 2.57 [1.37-4.4]; SIR for tracheal, bronchial, and lung cancers for employees in maintenance: 1.94 [1.03-3.31]). Negri et al. (Negri et al., 1989) found an association between bladder cancer and working in various departments (SMR for bladder cancer for employees in various departments: 3.5, p<0.05). Wang et al. (Wang et al., 1984) found an association between lung cancer and working in mixing and milling (SMR for lung cancer for employees in mixing and milling in the first company: 2.73, p<0.05; SMR for lung cancer for employees in mixing and milling in the second company 5.56, p<0.05).

De Vocht et al. (De Vocht et al., 2009) did not find any significant association despite using a matrix of exposure to aromatic amines except for the 3rd quartile of exposure (RR of bladder cancer for the third quartile of exposure to aromatic amines : 7,3 [1.1-51]).

Eleven studies have examined the overall risk of cancer. Of the nine studies that examined respiratory cancer, (Beall et al., 2007; Bernardinelli et al., 1987; McMichael et al., 1974; Mirabelli et al., 2012; Negri et al., 1989; Pira et al., 2012; Rubin & Clapp, 2012; Wilczyńska et al., 2001; Wingren & Axelson, 2007), two found an increased risk (McMichael et al., 1974; Wingren & Axelson, 2007). For bladder cancer, two (Negri et al., 1989; Veys, 2004) of the ten studies found an increased risk (Beall et al., 2007; Bernardinelli et al., 1987; McMichael et al., 1974; Mirabelli et al., 2012; Negri et al., 1989; Pira et al., 2012; Rubin & Clapp, 2012; Veys, 2004; Wilczyńska et al., 2001; Wingren & Axelson, 2007). For hematologic diseases and stomach cancer, only the McMichael et al. (McMichael et al., 1974) study found a significant increase in risk among the nine studies investigating it (Beall et al., 2007; Bernardinelli et al., 2012; Negri et al., 1989; Pira et al., 2007). For hematologic diseases and stomach cancer, only the McMichael et al. (McMichael et al., 1974) study found a significant increase in risk among the nine studies investigating it (Beall et al., 2007; Bernardinelli et al., 2012; Rubin & Clapp, 2012; Rubin & Axelson, 2007).

Among the five studies conducted in companies that were not all in the rubber tire sector (Boniol et al., 2016; Chen et al., 1997; McElvenny et al., 2018; Sorahan et al., 1989; Vlaanderen et al., 2013), two studies (McElvenny et al., 2018; Sorahan et al., 1989) found increased mortality from lung cancer, one study from stomach cancer (Chen et al., 1997), and one from bladder cancer (McElvenny et al., 2018) taking into account that Chen et al. (Chen et al., 1997) studied only digestive cancer.

## 4 **Discussion**

We found an increased risk of lung/respiratory cancer associated with a history of having worked in the tire industry across this literature review for several conditions of work.

This is the case for four of the twelve studies investigating overall increased risk or respiratory/lung cancer (McElvenny et al., 2018; McMichael et al., 1974; Sorahan et al., 1989; Wingren & Axelson, 2007). It is of note that for these studies the time periods of exposure are older than those of the studies that did not find any overall associated risk. Manufacturing processes have been improving over the years in the tire manufacturing industry, and carcinogen exposure as well. This result does not surprise us. Although the two studies that performed a analysis by hiring period, they did not find any association with cancer (Negri et al., 1989; Pira et al., 2012), it is possible that exposures in the oldest time periods will have been associated with an increased risk of lung/respiratory cancers. These data are consistent with data on occupational exposure to carcinogens, which show a decline in the intensity of exposure between the 1950s and 1990s in western countries (Boniol et al., 2016; Goutet, 2007).

On the other hand, through the analysis by work area, it appeared that some work areas are associated with an increased risk of developing cancer, specifically compounding, mixing, milling, and for employees who have cross-functional activities, whether in maintenance or in the various departments. Concerning compounding, mixing, and milling, these results are quite logical, as they are certainly the areas most exposed to carcinogens. The fumes released by the heating of raw materials contain PAHs, for example. However, these results should be taken with caution because the methods of classification of the employees into the different work areas are poorly described in the articles and certainly differ. Another work area that could have been found to be associated with some types of cancer is vulcanization, as exposure to fumes are also elevated in the area, but no significant association was found in the different analyses.

The link appears to be weaker for bladder cancer. Only three studies of the sixteen that studied the overall risk found an association (McElvenny et al., 2018; Negri et al., 1989; Veys, 2004). The studies of McElvenny et al. and Veys et al. are considered to hav a low risk of overall bias, but the study of Negri is considered to have a critical risk due to multiple analyses and it is thus impossible to conclude which work area could be more at risk. Exposure to aromatic amines is certain in the tire industry. Nvertheless, in the Polish cohort (De Vocht et al., 2009; Wilczyńska et al., 2001) no significant trend was found in the association between the indicator used to estimate exposure to aromatic amines and bladder cancer. However, the study of Veys et al. on

employees exposed to beta naphthylamine before 1949 is in favour of this association. The cessation of the use of beta naphthylamine occurred relatively early in comparison to the time periods examined by the studies in this review (1950s), and this may explain why the association seems weaker. Since then, other aromatic amines have replaced this substance. Concerning stomach cancer, the review does not provide argument supporting an association. Indeed, only two studies found a statistically significant association (Chen et al., 1997; McMichael et al., 1974). One had an overall risk of bias considered serious (Chen et al., 1997) among all the sixteen studies that examined this risk.

With regard to haematological malignancies, the classifications are not comparable between all the studies and the level of precision in the diagnosis is not identical, which makes it impossible to conclude on any precise trend. If we consider hematological diseases overall, only one study of sixteen found a statistically significant association (McMichael et al., 1974).

The analyses by duration of employment, time since first hired, time since end of exposure, age when hired, or time period of employment do not allow us to conclude that there is a trend between these indicators used to estimate exposure in terms of intensity or duration and the appearance of cancer.

Although the exposures are more homogeneous in the rubber tire industry than in the rubber industry more widely, an important limitation of the assemblage of the different studies is the fact that the level of protection and ventilation in the different companies studied, and even in the different work areas, is certainly not comparable.

There is great heterogeneity in the quality of the articles with regard to the assessment of the risk of bias, with two articles classified as critical. The majority of the studies that found statistically significant associations had a risk of bias assessment classified between low and moderate (McElvenny et al., 2018; McMichael et al., 1974; Sorahan et al., 1989; Veys, 2004; Wingren & Axelson, 2007), which supports our interpretation. However, it should be noted that only one of the authors assessed the risk of bias.

Our systematic review initially relied on the MEDLINE database because it is more accessible to algorithmic searching. However, we cross-referenced our data with the references of the articles, with those of the meta-analyses cited previously (Alder et al., 2006; Boniol et al., 2017) and a search on WEB OF SCIENCE. Two articles emerged from this cross-referencing, which suggests good sensitivity of our method, especially since the overall risk of bias is considered serious (Chen et al., 1997) or critical (Rubin & Clapp, 2012) for these articles.

## 5 Conclusion

This systematic review has highlighted that, despite the results of meta-analyses that did not find any excess risk of cancer in the rubber tire industry (Alder et al., 2006; Boniol et al., 2017), having worked in this industry, especially in older time periods likely having higher exposure to carcinogens, could be associated with an increased risk of lung cancer. This is particularly true for specific jobs exposing workers to fumes. However, there is no evidence to suggest that this risk is proportional to the duration of exposure.

Future studies on the tire industry should take into account, at minimum, the smoking status of àrthe employees, and at best the major exposures to carcinogens by work area. All the studies reviewed here lacked precision on these points, which limits the conclusions we can draw from the trends described.

#### Declarations

The authors did not receive support from any organization for the submitted work.

All selected articles are available on the internet and data on their selection and evaluation are available in the supplementary data of the manuscript.

The first author wrote the manuscript, selected the articles and assessed their risk of bias. The second author selected the articles with the first author and corrected the manuscript. The other authors corrected the manuscript.

### 6 <u>References</u>

- Alder, N., Fenty, J., Warren, F., Sutton, A. J., Rushton, L., Jones, D. R., & Abrams, K. R. (2006). Meta-analysis of mortality and cancer incidence among workers in the synthetic rubber-producing industry. *American Journal of Epidemiology*, 164(5), 405-420. https://doi.org/10.1093/aje/kwj252
- Beall, C., Corn, M., Cheng, H., Matthews, R., & Delzell, E. (2007). Mortality and cancer incidence among tire manufacturing workers hired in or after 1962. *Journal of Occupational and Environmental Medicine*, 49(6), 680-690. https://doi.org/10.1097/JOM.0b013e318074bb30
- Bernardinelli, L., de Marco, R., & Tinelli, C. (1987). Cancer mortality in an Italian rubber factory. *British Journal of Industrial Medicine*, 44(3), 187-191.
- Boniol, M., Koechlin, A., & Boyle, P. (2017). Meta-analysis of occupational exposures in the rubber manufacturing industry and risk of cancer. *International Journal of Epidemiology*. https://doi.org/10.1093/ije/dyx146
- Boniol, M., Koechlin, A., Świątkowska, B., Sorahan, T., Wellmann, J., Taeger, D., Jakobsson, K., Pira, E., Boffetta, P., La Vecchia, C., Pizot, C., & Boyle, P. (2016). Cancer mortality in cohorts of workers in the European rubber manufacturing industry first employed since 1975. *Annals of Oncology: Official Journal of the European Society for Medical Oncology*, 27(5), 933-941. https://doi.org/10.1093/annonc/mdw061
- Chen, J., Wei, X., & You, X. (1997). A retrospective cohort study on digestive cancer in the rubber tire industry in Shangai. *Journal of Occupational Health*, *39*, 302-312.
- Coggon, D., Barker, D. J., & Cole, R. B. (1990). Stomach cancer and work in dusty industries.
   *Occupational and Environmental Medicine*, 47(5), 298-301.
   https://doi.org/10.1136/oem.47.5.298

- De Vocht, F., Sobala, W., Wilczynska, U., Kromhout, H., Szeszenia-Dabrowska, N., & Peplonska, B. (2009). Cancer mortality and occupational exposure to aromatic amines and inhalable aerosols in rubber tire manufacturing in Poland. *Cancer Epidemiology*, 33(2), 94-102. https://doi.org/10.1016/j.canep.2009.06.013
- Dost, A. A., Redman, D., & Cox, G. (2000). Exposure to rubber fume and rubber process dust in the general rubber goods, tyre manufacturing and retread industries. *The Annals of Occupational Hygiene*, 44(5), 329-342.
- Fox, A. J., & Collier, P. F. (1976). A survey of occupational cancer in the rubber and cablemaking industries : Analysis of deaths occurring in 1972-74. Occupational and Environmental Medicine, 33(4), 249-264.
- Goutet, P. (2007). Evolution des expositions industrielles aux cancérogènes vésicaux. 112, 533-536.
- Grossetta Nardini, H., & Wang, L. (2022). The Yale MeSH Analyzer [Internet]. *CT: Cushing/Whitney Medical Library*. http://mesh.med.yale.edu/
- Héry, M. (2009). *Cancers de la vessie et risques professionnels*. INRS, Institut national de recherche et de sécurité EDP sciences.
- Hidajat, M., McElvenny, D. M., Mueller, W., Ritchie, P., Cherrie, J. W., Darnton, A., Agius,
  R. M., Kromhout, H., & de Vocht, F. (2019). Job-exposure matrix for historical exposures to rubber dust, rubber fumes and n-Nitrosamines in the British rubber industry. *Occupational and Environmental Medicine*, 76(4), 259-267. https://doi.org/10.1136/oemed-2018-105182
- International Agency for Research on Cancer. (1982). *IARC monographs on the evaluation of carcinogenic risks to humans 28 : The Rubber Industry*. International Agency for Research on Cancer.

- International Agency for Research on Cancer. (1987). Overall evaluations of carcinogenicity : An updating of IARC monographs volumes 1 to 42. International Agency for Research on Cancer.
- International Agency for Research on Cancer. (2012). *IARC monographs on the evaluation of carcinogenic risks to humans 100F : A review of human carcinogens*. International agency for research on cancer.
- McElvenny, D. M., Mueller, W., Ritchie, P., Cherrie, J. W., Hidajat, M., Darnton, A. J., Agius,
  R. M., & de Vocht, F. (2018). British rubber and cable industry cohort: 49-year mortality follow-up. *Occupational and Environmental Medicine*, 75(12), 848-855. https://doi.org/10.1136/oemed-2017-104834
- McMichael, A. J., Spirtas, R., & Kupper, L. L. (1974). An epidemiologic study of mortality within a cohort of rubber workers, 1964-72. *Journal of Occupational Medicine.: Official Publication of the Industrial Medical Association*, 16(7), 458-464.
- Mirabelli, D., Cacciatore, A. M., Ferrante, D., Amendola, P., Vermeulen, R., & Merletti, F. (2012). Cohort study of workers employed in an Italian tire manufacturing plant, 1962-2004. *Cancer Causes & Control: CCC*, 23(12), 2023-2029. https://doi.org/10.1007/s10552-012-0083-y
- Negri, E., Piolatto, G., Pira, E., Decarli, A., Kaldor, J., & La Vecchia, C. (1989). Cancer mortality in a northern Italian cohort of rubber workers. *British Journal of Industrial Medicine*, 46(9), 624-628.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., ... Moher, D. (2021). The PRISMA 2020 statement : An updated

guideline for reporting systematic reviews. *BMJ (Clinical Research Ed.)*, 372, n71. https://doi.org/10.1136/bmj.n71

- Pira, E., Pelucchi, C., Romano, C., Manzari, M., Negri, E., & La Vecchia, C. (2012). Mortality from cancer and other causes in an Italian cohort of male rubber tire workers. *Journal* of Occupational and Environmental Medicine, 54(3), 345-349. https://doi.org/10.1097/JOM.0b013e318244503c
- ROBINS-E Development Group. (2022). Risk Of Bias In Non-randomized Studies—Of Exposure (ROBINS-E). https://www.riskofbias.info/welcome/robins-e-tool
- Rubin, S. M., & Clapp, R. (2012). Patterns of mortality among Wisconsin Uniroyal Tire Manufacturing workers. NEW SOLUTIONS: A Journal of Environmental and Occupational Health Policy, 21(4), 603-620. https://doi.org/10.2190/NS.21.4.h
- Sorahan, T., Parkes, H. G., Veys, C. A., Waterhouse, J. A., Straughan, J. K., & Nutt, A. (1989).
  Mortality in the British rubber industry 1946-85. *British Journal of Industrial Medicine*, 46(1), 1-10.
- Veys, C. A. (2004). Bladder tumours in rubber workers: A factory study 1946–1995. Occupational Medicine, 54(5), 322-329. https://doi.org/10.1093/occmed/kqh010
- Vlaanderen, J., Taeger, D., Wellman, J., Keil, U., Schüz, J., & Straif, K. (2013). Extended cancer mortality follow-up of a German rubber industry cohort. *Journal of Occupational and Environmental Medicine*, 55(8), 966-972. https://doi.org/10.1097/JOM.0b013e31829540f4
- Wang, H., You, X., & Qu, Y. (1984). Investigation of Cancer Epidemiology and Study of Carcinogenic Agents in the Shanghai Rubber Industry. 7.
- Weiland, S. K., Straif, K., Chambless, L., Werner, B., Mundt, K. A., Bucher, A., Birk, T., &Keil, U. (1998). Workplace risk factors for cancer in the German rubber industry : Part

1. Mortality from respiratory cancers. *Occupational and Environmental Medicine*, 55(5), 317-324.

- Wilczyńska, U., Szadkowska-Stańczyk, I., Szeszenia-Dabrowska, N., Sobala, W., & Strzelecka, A. (2001). Cancer mortality in rubber tire workers in Poland. *International Journal of Occupational Medicine and Environmental Health*, 14(2), 115-125.
- Wingren, G., & Axelson, O. (2007). Cancer incidence and mortality in a Swedish rubber tire manufacturing plant. American Journal of Industrial Medicine, 50(12), 901-909. https://doi.org/10.1002/ajim.20536

Figure 1. Flow chart of literature search for observational studies on occupational exposure in the tire manufacturing industry and risk of cancer.



**Table 1. Summary of studies from the literature review on occupational exposures in the tire industry and risk of cancer.**Attached to the article.

Country/Ref
USA/Beall et al.
(2007)
(2007)

Only case-con- trol study in the rubber the review, tire indust concerning 190 age ar stomach can- cer only controls	Comparison to lo- cal population Taking into ac- count the diet Questionnaire and interview about diet and occupa- tion	Newly diag- nosed case during 1985- 1987, no other infor- mation	RR for stomach cancer for employ- ees exposed 5 years or more at least ten year be- fore the study RR for stomach cancer for employ- ees exposed less than 5 years or shorter than ten year before the study	3.5 [1.1-11] 0.9 [0.2-4.3]	RR for stomach cancer taking acount of diet	2.5 [1-6.4]	England/Coggon et al. (1990) et al. (1997)
1 tire manu- facturing plant (Overlap with Boniol 2016 (36.4% of workers hired after 1975)	Vital status known for 99% of employ- ees, cause of death in 96% of cases Comparison to lo- cal population (Piedmont) 10 year lag	1962-2000 1962-2004	Multivariate re- gression ratio (Poisson) for blad- der cancer inci- dence for employ- ees exposed from 5 to 9 years Multivariate re- gression ratio (Poisson) for blad- der cancer inci- dence for employ- ees exposed from 10 to 19 years Multivariate re- gression ratio (Poisson) for blad- der cancer inci- dence for employ- ees exposed more than 20 years	1.03 [0.4-2.8] 1.57 [0.7-3.3] 1,74 [0,9-3,6]	SIR for bladder cancer in males	1.15 [0.9-1.4]	Italy/Mirabelli et al. (2012)

facturing plant	employee has worked less than 12 months	1946-1981	cer for employees exposed more than 20 years	0.97	SMR for lung cancer SMR for lung cancer for employees in various services <sup>b</sup>	1.01 [0.79-1.29] 2.25*	(1989)
	Comparison to na- tional population (Italy) Analyse by du- rartion of expo- sure, time since last exposure, age				SMR for lung cancer for employees hired be- tween 1906 and 1939 SMR for bladder cancer SMR for bladder cancer for employees in various services <sup>b</sup>	1.27 1.83 [1.05-2.96] 3.5*	
	at first exposure, time period at first employment and by job title				SMR for stomach SMR for leukemia	0.78 [0.54-1.08] 0.91 [0.39-1.79]	

1 tire manu- facturing plant	6,246	Exclusion if the employee has worked less than 3 months or is over	1954-2008 1954-2008	SMR for bladder cancer for employ- ees exposed less than 10 years	0.81 [0.29-1.76]	SMR for lung cancer SMR for lung cancer for employees hired before 1965	0.8 [0.6-0.9] 0.7 [0.57-0.97]	Italy/Pira et al. (2012)
		worked less than 3 months or is over 85 years old Vital status known for 95% of employ- ees, cause of death in 94% of cases Comparison to lo- cal population (Piedmont) Analyse by du- rartion of employ- ment, time since first employment, time since last em- ployment and time period at first em- ployment		ees exposed less than 10 years SMR for bladder cancer for employ- ees exposed from 10 to 19 years SMR for bladder cancer for employ- ees exposed more than 20 years	1.83 [0.79-3.6] 1.27 [0.55-2.51]	employees hired before 1965 SMR for stomach cancer SMR for all lymphatic and hematopoetic can- cers	1 [0.7-1.4] 1 [0.7-1.4]	

1 tire manu-	17,636	Exclusion if the	1950-1995	SMR for bladder	0.56 [0.07-2.02]	SMR for lung cancer	0.7 [0.6-0.9]	Poland/ de Vocht et
facturing		employee has	1950-2001	cancer for employ-		SMR for bladder in	0.6 [0.2-1.3]	al. (2009) and
plant, Overlap		worked less than 3		ees exposed less		males		Wilczyńska et al.
with Boniol		months		than 5 years		SMR for bladder in	1.3 [0.3-3.8]	(2001)
2016 (~60% of		Vital status known		SMR for bladder	0.64 [0.17-1.64]	males in preparation of		
workers hired		for 97% and cause		cancer for employ-		materials <sup>b</sup>		
after 1970)		of death for 88%.		ees exposed more		RR of bladder cancer for	7,3 [1.1-51]	
		Comparison to na-		than 5 years		the third quartile of ex-		
		tional population				posure to aromatic		
		(Poland)				amines		
		Use Exposure Ma-				RR of bladder cancer for	8.3 [1-66.3]	
		trix for aromatic				the fourth quartile of		
		amines and inhala-				exposure to aromatic		
		ble aerosols				amines		
		Latency period of 5				SMR for tumors of lym-	0.6 [0.4-1.1]	
		years				phoid and hematopoi-		
						etic tissues in males		
						RR for tumors of lym-	2.1 [0.5-7.9]	
						phoid and hematopoi-		
						etic tissues in males for		
						the second quartile of		
						exposure to aromatic		
						amines		
						RR for tumors of lym-	1.3 [0.2-7.2]	
						phoid and hematopoi-		
						etic tissues in males for		
						the third quartile of ex-		
						posure to aromatic		
						amines		
						RR for tumors of lym-	3 [0.8-11.1]	
						phoid and hematopoi-		
						etic tissues in males for		
						the fourth quartile of		
						exposure to aromatic		
						amines		

1 tire manu- facturing plant	5,745	Exclusion if the employee has worked less than 12 months Comparison to na- tional population	1938-1981 1958-2001	SIR for trachea, bronchus, and lung cancers for em- ployees exposed from 2,5 to 10 years	1.75 [1.04-2.77]	SIR for trachea, bron- chus, and lung cancers SMR for trachea, bron- chus, and lung cancers SIR for trachea, bron- chus, and lung cancers	1.62 [1.28-2.02] 1.54 [1.21-1.94] 2.18 [1.31-3.4]	Sweden/Wingren et al. (2007)
		(Sweden) Latency period of 10 years		SIR for trachea, bronchus, and lung cancers for em- ployees exposed more than 10	1.78 [1.09-2.75]	for employees in com- pounding and mixing <sup>b</sup> SIR for trachea, bron- chus, and lung cancers for employees in miling <sup>b</sup>	2.57 [1.37-4.4]	
				years		SIR for trachea, bron- chus, and lung cancers for employees in maintenance <sup>b</sup>	1.94 [1.03-3.31]	

2 tire manu- facturing plants	not find	Included if the employee is over 30 years old Comparison to lo- cal population (Shangai)	1971-1980 and 1976- 1980 (differ- ence be- tween expo- sure and fol- low-up peri- ods not clear)		SMR for lung cancer for employees in mixing and milling in the first plant SMR for lung cancer for employees in mixing and milling in the sec- ond plant <sup>b</sup> SMR for stomach cancer for employees in mixing and milling in the first plant <sup>b</sup> SMR for stomach cancer for employees in mixing and milling in the sec- ond plant <sup>b</sup>	2.73* 5.56* 1.59 1.84	China/Wang et al. (1984)
1 tire manu- facturing plant	4,917	Exclusion if the employee has worked less than 12 months Vital status known for 99% of employ- ees Comparison to lo- cal population (Cu- neo province) Only males	1962-1983 1962-1983		SMR for stomach cancer SMR for bronchus, tra- chea or lung cancers SMR for bladder cancer SMR for Hodgkin lym- phoma SMR for myeloid leuke- mia SMR for other leukemia	1.4 0.7 1.4 1.7 3.6 1.1	ltaly/Bernardinelli et al. (1987)

1 tire manu- facturing plant	6,678	Inclusion of active and retired em- ployees aged 40 to 84 at 1/1/1964 Vital status known for 99% of employ- ees, cause of death in 99.5% of cases Only male	Until 1972 (no infor- mation on the opening date of the plant) 1964-1972		SMR for stomach cancer SMR for respiratory sys- tem cancer SMR for bladder cancer SMR for lymphosarcoma SMR for leukemia	1.87* 0.84* 0.73 2.26* 1.28	USA/Mc Michael et al. (1974)
1 tire manu- facturing plant	5,128	Vital status known for 99,8% of em- ployees Comparison to lo- cal national (Eng- land and Wales) and local (Birming- ham) population	1945-1949 1951-1995		SIR for bladder SIR for bladder for em- ployees exposed to beta-naphtylamine be- tween 1946 and 1949 SMR for bladder SMR for bladder for em- ployees exposed to beta-naphtylamine be- tween 1946 and 1949	1.23 [1.02-1.48] 1.71 [1.3-2.21] 0.97 [0.55-1.57] 0.85 [0.6-1.18]	Great Britain/Veys et al. (2004)

1 tire manu-	2,924	Exclusion if the	1932-1992		PCMR for lung cancer in	0.9 [0.64-1.22]	USA/Rubin et al.
facturing plant		employee has	No informa-		males		(2012)
		worked less than	tion		PCMR for bladder can-	0.46 [0.06-1.67]	
		12 months			cer in males		
		Comparison to lo-			PCMR for stomach can-	1.21 [0.39-2.82]	
		cal population			cer in males		
		(Winsconsin)			PCMR for leukemia can-	0.98 [0.36-2.13]	
		Latency period of 5			cer in males		
		years			PCMR for leukemia can-	3.32 [0.68-9.7]	
					cer in males hired be-		
					twenn 1932 and 1950		
1							

Rubber in-	17,389 in	Pooled analysis of	1975-(max		SMR for lung cancer in	0.71 [0.56-0.90]	Germany, Italy, Po-
dustry, 64	tire manu-	5 European co-	2013)1975-		tire plants		land, Sweden and
plants	facturing	horts	(max 2013)		SMR for bladder cancer	0.97 [0.46-2.08]	Great Britain/Boniol
	plants	Exclusion if the			in tire plants		et al. (2016)
		employee has			SMR for stomach cancer	0.92 [0.59-1.44]	
		worked less than			in tire plants		
		12 months			SMR for leukemia can-	0.96 [0.52-1.78]	
		Only first em-			cer in tire plants		
		ployed at 1 January					
		1975 or later					
		Comparison with					
		national popula-					
		tion according to					
		WHO data, except					
		in Italy					
		Overlap with de					
		Vocht et al. (2009)					
		and Wilczyńska et					
		al. (2001) with &					
		Mirabelli et al.					
		(2012)					
		· · /					

3 tire and other rubber goods manu- facturing plants	7,808, num- ber of em- ployees working in tire manu- facturing not specified	Exclusion if the employee has worked less than 12 months Comparison to lo- cal population (Shanghai) Taking into ac- count the smoking status Latency period of 5 years	1920s (max) -1972 1967-1988		SIR for stomach cancer in non-smokers males in miling and smelting for tire SMR for stomach cancer in males in material handling for tire	2.3* 2*	China/Chen et al. (1997)
Rubber in- dustry	14,906 in tire manu- facturing plants	Employee younger than 35 years of age were excluded Comparison to lo- cal population (England and Wales)	Recruitment in 1967 but could have worked since 1950 or before 1967-2015		SMR for trachea, bron- chus, and lung cancers in tire plants SMR for bladder cancer in tire plants	1.25 [1.19-1.32] 1.38 [1.23-1.53]	Great Britain/McEl- venny et al. (2018)

Rubber in- dustry, 13 plants	36,445, number of employees	Exclusion if the employee has worked less than	1946-1960 1946-1985		SMR for stomach cancer for employees hired be- tween 1946 and 1950	1.12	Great Britain/Sora- han et al. (1989)
F	working in	12 months Vital status known			SMR for stomach cancer	0.97	
	facturing not	for 93% of employ-			tween 1951 and 1955		
	specified	ees			SMR for stomach cancer	1.29	
		Comparison to lo-			for employees hired be-		
		cal population			tween 1956 and 1960		
		(England and			SMR for lung cancer for	1.32	
		Wales)			employees hired be-		
					SMR for lung cancer for	1 16	
					employees hired be-	1.10	
					tween 1951 and 1955		
					SMR for lung cancer for	1.33	
					employees hired be-		
					tween 1956 and 1960		

Rubber in-	13,495 in these 5	Exclusion if the	Recruitment		SMR for lung cancer in production of tire	1.09 [0.87-1.36]	Germany/Vlaande-
plants	plants	worked less than	could have		SMR for bladder cancer	1.31 [0.7-2.23]	10110101. (2013)
		12 months	worked be-		in production of tire		
		No analyse by du-	fore		SMR for stomach cancer	1.17 [0,75-1.72]	
		ration of employ-	1981-2000		in production of tire		
		ment in tire indus-			SMR for leukemia can-	0.39 [0.08-1.15]	
		try			cer in production of tire		
		Comparison with					
		local population					
		(Western Ger-					
		many)					
		Vital status known					
		for 99% of employ-					
		ees, , cause of					
		death in 93% of					
		cases (males)					
		work by duration of					
		employment and					
		time since first em-					
		nlovment					
		ployment					

\* p<0,05</li>
 <sup>a</sup> only for a priori important causes of death based on the IARC monograph conclusions for rubber industry
 <sup>b</sup> analysis by work area

First author (year)	D1	D2	D3	D5	D6	D7	Overall
McElvenny (2018)							
Boniol (2016)							
Vlaanderen (2013)							
Mirabelli (2012)							
Pira (2012)							
de Vocht (2009)							
Wingren (2007)							
Veys (2004)							
Beall (2007)							
Negri (1989)							
Sorahan (1989)							
Bernardinelli (1987)							
Wang (1984)							
Mc Michael (1974)							
Chen (1997)							
Coggon (1990)							
Rubin & Clapp (2012)							

Figure 2. Risk of bias summary : reviewers' judgement regarding each risk of bias domain foe each included study.

Domains: D1: Bias due to confounding, D2: Bias in measurement of the exposure, D3: Bias in selection of participants into the study, D5: Bias due to missing data, D6: Bias in measurement of outcomes and D7: Bias in the selection of the reported results.

Judgement:

