



HAL
open science

Influence of cell wall pectic fraction modification on Cadmium stress capacities

Jonathan Robilliard, Sophie Bouton, Karine Pageau, Romain Roulard,
Stéphanie Guénin, Laurent Gutierrez, Pierre Delaplace, Olivier van
Wuytswinkel

► **To cite this version:**

Jonathan Robilliard, Sophie Bouton, Karine Pageau, Romain Roulard, Stéphanie Guénin, et al..
Influence of cell wall pectic fraction modification on Cadmium stress capacities. XVI plant cell wall
meeting, Jun 2023, Malaga, Espagne, Spain. 10.1016/j.ijbiomac.2021.02.035.;[9] . hal-04425948

HAL Id: hal-04425948

<https://u-picardie.hal.science/hal-04425948>

Submitted on 30 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Jonathan Robilliard^{1,3}, Sophie Bouton¹, Karine Pageau¹, Romain Roulard¹, Stéphanie Guénin², Laurent Gutierrez², Pierre Delaplace³, Olivier Van Wuytswinkel¹

¹Université de Picardie Jules Verne - UMRT INRAE 1158 BioEcoAgro - Biologie des Plantes et Innovation (BioPI)

² Université de Picardie Jules Verne – Centre de Ressources Régionales en Biologie Moléculaire (CRRBM)

³ Université de Liège - Gembloux Agro-Bio Tech (GxABT) – TERRA - Plant Sciences

Abstract


Pectin MethylEsterases (PME) catalyze the hydrolysis of the C6 Galacturonic Acid (GalA) methylester group present in newly synthesized HomoGalacturonans (HG) [1]. Analysis of recent literature suggests a potential role for pectin demethylesterification in the plant's response to nutritional stresses such as nitrogen and phosphorus [2, 3, 4] or metal stresses such as cadmium and aluminum [5, 6, 7]. This is evidenced by changes in PME activity under these conditions. These observations raise the question of whether pectin methylesterification rate is involved in the apoplastic transport and translocation of ions in both root and aerial plant tissues.

Objective


Assess how modifications to the cell wall pectic fraction impact the absorption and translocation of cadmium within *Arabidopsis thaliana*

Results

1 Impact of modifications on the absorption and transport of Cadmium



3 Accumulation of cadmium in leaves



2 Characterization of *A. thaliana* cell wall mutants


Double mutant *gaut10gaut11*
(GalactUronosyl Transferase → synthesis of HG)

- ∅ visible phenotypic difference between *gaut10gaut11* and Col 0
- Lower amount of GalA in cell wall (obtained by F. Sénéchal and P. Trezlet : BioEcoAgro)


Double mutant *cgr2cgr3*
(HG MethylTransferase → HG methylesterification)

- Growth ∇ (Dwarf phenotype)
- Early appearance of inflorescence
- Defect in silique formation
- Lower methylesterification rate (Kim et al. 2015)

Obtained thanks to F. Brandizzi



4 Effect of Cadmium on cell wall composition



Conclusions

- gaut10gaut11* mutant did show a significant decrease of GalA in addition to a decrease of Rhamnose
- Cadmium stress induces a decrease in GalA content in roots cell wall
- Metallic ions accumulation in *cgr2cgr3* leaves, whatever condition, suggests that lower methylesterification level results in cation accumulation
- Upon cadmium stress, accumulation of cadmium in *cgr2cgr3* mutants is slightly decreased. Which can be explained by an overload of cell wall by other ions as shown by copper and iron hyperaccumulation

Perspectives

- Plants overexpressing a fungal PME [8] in different tissues will be tested to confirm importance of methylesterification rate under cadmium stress conditions
- Cadmium accumulation kinetics and the effect of cadmium on cell wall modifications will be tested
- Influence of cadmium on root architecture will be tested on different mutants
- Capacity of cell wall to bind cadmium will be tested using extracted cell wall from different mutants
- PME and PMEi genes expression upon cadmium stress will be measured using RT-qPCR

References

- [1] Sénéchal et al. (2014) J Exp Bot 65:5125. ; [2] Landi S. and Esposito S. (2017) Front. Plant Sci. 8: 1376. ; [3] Podgorska et al. (2017) Front. Plant Sci. 8: 1344. ; [4] Hoehenwarter et al. (2015) BMC Plant Biol 16: 106. ; [5] Li et al. (2017) Plant Physiology and Biochemistry 119: 265. ; [6] Corso et al. (2021) New Phytol. 230: 669. ; [7] Wu et al. (2021) Environmental Pollution doi.org/10.1016/j.envpol.2021.117218 ; [8] Safran J et al. (2021) Int J Biol Macromol. 15:176:165-176. doi: 10.1016/j.ijbiomac.2021.02.035. ; [9] Schürholz AK et al. (2018) Plant Physiol. 2018 Sep;178(1):40-53. doi: 10.1104/pp.18.00463. Epub 2018 Jul 19.