



HAL
open science

Sugar-Based Surfactants: Effects of Structural Features on the Physicochemical Properties of Sugar Esters and Their Comparison to Commercial Octyl Glycosides

Huiling Lu, Gwladys Pourceau, Benoit Briou, Anne Wadouachi, Théophile Gaudin, Isabelle Pezron, Audrey Drelich

► **To cite this version:**

Huiling Lu, Gwladys Pourceau, Benoit Briou, Anne Wadouachi, Théophile Gaudin, et al.. Sugar-Based Surfactants: Effects of Structural Features on the Physicochemical Properties of Sugar Esters and Their Comparison to Commercial Octyl Glycosides. *Molecules*, 2024, 29 (10), pp.2338. 10.3390/molecules29102338 . hal-04580210

HAL Id: hal-04580210

<https://u-picardie.hal.science/hal-04580210v1>

Submitted on 24 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Article

Sugar-Based Surfactants: Effects of Structural Features on the Physicochemical Properties of Sugar Esters and Their Comparison to Commercial Octyl Glycosides

Huiling Lu ¹, Gwladys Pourceau ^{2,*}, Benoit Briou ², Anne Wadouachi ², Théophile Gaudin ¹,
Isabelle Pezron ¹ and Audrey Drelich ^{1,*}

¹ Université de Technologie de Compiègne, ESCOM, TIMR (Transformations Intégrées de la Matière Renouvelable), Centre de Recherche Royallieu—CS 60 319, 60203 Compiègne Cedex, France; lhl@ujs.edu.cn (H.L.); isabelle.pezron@utc.fr (I.P.)

² Laboratoire de Glycochimie, et des Agroressources d'Amiens (LG2A), UR 7378—Institut de Chimie de Picardie, Université de Picardie Jules Verne, 33 rue Saint Leu, 80039 Amiens Cedex, France; b.briou@orpiainnovation.com (B.B.); anne.wadouachi@u-picardie.fr (A.W.)

* Correspondence: gwladys.pourceau@u-picardie.fr (G.P.); audrey.drelich@utc.fr (A.D.)

Abstract: Two series of sugar esters with alkyl chain lengths varying from 5 to 12 carbon atoms, and with a head group consisting of glucose or galactose moieties, were synthesized. Equilibrium surface tension isotherms were measured, yielding critical micellar concentration (CMC) surface tensions at CMC (γ_{cmc}) and minimum areas at the air–water interface (A_{min}). In addition, Krafft temperatures (T_k) were measured to characterize the ability of molecules to dissolve in water, which is essential in numerous applications. As a comparison to widely used commercial sugar-based surfactants, those measurements were also carried out for four octyl D-glycosides. Impacts of the linkages between polar and lipophilic moieties, alkyl chain lengths, and the nature of the sugar head group on the measured properties were highlighted. Higher T_k and, thus, lower dissolution ability, were found for methyl 6-O-acyl-D-glucopyranosides. CMC and γ_{cmc} decreased with the alkyl chain lengths in both cases, but A_{min} did not appear to be influenced. Both γ_{cmc} and A_{min} appeared independent of the ester group orientation. Notably, alkyl (methyl α -D-glucopyranosid)uronates were found to result in noticeably lower CMC, possibly due to a closer distance between the carbonyl function and the head group.

Keywords: sugar esters; carbohydrate-based surfactants; glycolipids; physicochemical parameters



Citation: Lu, H.; Pourceau, G.; Briou, B.; Wadouachi, A.; Gaudin, T.; Pezron, I.; Drelich, A. Sugar-Based Surfactants: Effects of Structural Features on the Physicochemical Properties of Sugar Esters and Their Comparison to Commercial Octyl Glycosides. *Molecules* **2024**, *29*, 2338. <https://doi.org/10.3390/molecules29102338>

Academic Editors: Khristo Khristov and Plamen Tchoukov

Received: 22 March 2024

Revised: 9 May 2024

Accepted: 10 May 2024

Published: 16 May 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Glycolipids are often considered safe alternatives to petroleum-based surfactants due to their renewable origin, high biodegradability, and low toxicity. Their surface-active properties make them highly interesting in a wide range of fields such as food, cosmetics, pharmaceuticals, and detergent industries [1–5]. These amphiphilic molecules comprise a saccharidic polar head group linked to a lipophilic chain. Depending on the linkage, its orientation and position, the length and degree of saturation of the alkyl chain, the sugar residue and its size, etc., a wide variety of sugar-based amphiphiles can be designed. Despite the fact that many studies have focused on relationships between their chemical structures and surface properties, important gaps remain, owing to the structural versatility of sugar-based surfactants [6–9]. Filling those gaps (regarding the knowledge of structure–property relationships) will help identify and anticipate trends, with the final aim of designing green bio-based surfactants for specific applications.

The structures of sugar-based surfactants are diverse (including glyco-amides, glyco-esters, glycosides, and other derivatives), and their properties also widely vary. For example, in previous work [10], we synthesized various glyco-amides, bearing an octyl chain, which

differed by the position and the orientation of the amide linkage on the sugar head-group (Su-NHCO-R or Su-CONH-R), and measured significant differences for those molecules in physicochemical properties, such as solubility or CMC, as well as in cytotoxic effects. In particular, esterase enzymes found widely in nature can break down sugar esters, leading to high biodegradability for this specific surfactant family, and making it relevant whenever eco-compatibility is a concern [11]. Another advantage of sugar esters is the general ease of their synthesis, as the ester linkage can be obtained by simple chemical [12–14] or enzymatic routes [15–18]. However, their ability to dissolve in water appears to be lower than that of commercial alkyl glycosides, a difference observed across a diverse range of sugar residues combined with ester linkages (alkyl uronates, alkyl aldonates, acyl oses, acyl itols, etc.) [19–24]. Moreover, in most studies, mixtures of α and β isomers with an undetermined ratio amount are reported, even though this structural parameter influences dissolution and micellization [25–28]. As reported by Brown et al. [25], the solubility of octyl- β -D-glucoside (C8 β Glc) is about seven times larger than that of octyl- α -D-glucoside (C8 α Glc), due to the latter's more stable crystal structure. Consistently, glucosides with α -linkages are generally believed to have higher Krafft temperatures (Tks) than those with β -linkages [26]. Thus, it is essential to know exactly the α/β ratio or to protect the anomeric position to establish structure/function relationships. It is also necessary to keep in mind that the replacement of the anomeric -OH function with other groups should cause a decreased solubility of the corresponding esters at 25 °C [20]. In 2016, we reported on the synthesis and physicochemical behavior of one family of sugar esters with well-defined anomeric configurations, namely 1-O-methyl alkyl uronates (Me α Glc, Me β Gal, and Me α Man derivatives) (Figure 1 left) [29]. This study revealed that, similar to corresponding alkyl polyglycosides or common polyoxyethylene nonionic surfactants, our newly synthesized sugar esters were able to significantly reduce the surface tensions of aqueous solutions, at low concentrations. Nevertheless, their solubility remained limited.




Figure 1. Structures of 1-O-methyl alkyl-D-glycuronates [29] (left) and methyl 6-O-acyl-D-glycopyranosides (right). Each molecule is abbreviated with the code annotated below.

Thus, we synthesized and characterized close, but distinct sugar esters with well-defined anomeric positions, derived from Me α Glc and Me β Gal: methyl 6-O-acyl-D-glycopyranosides (Figure 1 right). Their physicochemical properties were compared to the corresponding 1-O-methyl alkyl uronates that we previously described. Thanks to the homology of both families (the same nature and position of the linkage, and the same anomeric configuration), the effects of other structural parameters, including the chain length (hexyl to dodecyl derivatives), the orientation of the ester groups, as well as the sugar head, can be directly compared.

Commercial alkyl α/β -glycosides have been, by far, the most extensively investigated sugar-based surfactants, making them relevant comparison models. Indeed, their

structures are a straightforward combination of a typical sugar residue, such as glucose, with an alkyl chain, i.e., a prototypical surfactant that can be imagined with sugar as a polar head. Nevertheless, confidence in the available knowledge of their physicochemical behavior is limited by significant inconsistency and discrepancy for the same structure sometimes apparent within a large amount of published physicochemical measurements, as we highlighted previously [30]. For example, in 1997, Sakya et al. showed—for the first time—the phase diagram of octyl- β -D-galactoside (C8 β Gal), from which a Tk of 42 °C was deduced by Hato et al. [31]. In parallel, several researchers have measured the surface tension of C8 β Gal directly at 25 °C, assuming that the surfactant properly dissolves in water at this temperature [32,33]. To clarify the observed discrepancies in the literature regarding the structure–property relationships for alkyl glycosides, in the present study, properties for two pairs of α / β anomers, i.e., C8 α Glc/C8 β Glc and C8 α Gal/C8 β Gal, have also been investigated (Figure 2).





Figure 2. Structures of octyl- α / β -D-glucosides (**left**) and octyl- α / β -D-galactosides (**right**). Each molecule is abbreviated with the code annotated below.

2. Results and Discussion

2.1. Synthesis

Two families of sugar esters, alkyl (methyl D-glycopyranosid)uronates, and methyl 6-*O*-acyl-D-glycopyranosides, were synthesized. The uronates family was obtained using the two-step procedure that we previously described [29]. For the methyl 6-*O*-acyl-D-glycopyranosides, a known strategy ensuring regioselectivity on OH-6 [34] was successfully adapted and applied to methyl D-glycosides (Scheme 1). Firstly, α -methyl D-glucoside was per-silylated by treating with *N,O*-bis(trimethylsilyl)acetamide (BSA) and catalytic tetrabutylammonium fluoride (TBAF), and the more labile primary silyl ether was deprotected in a one-pot procedure during the work-up by the addition of K_2CO_3 to generate methyl 2,3,4-tri-*O*-trimethylsilyl- α -D-glycopyranoside **1** with an 80% yield. Methyl 2,3,4-tri-*O*-trimethylsilyl- β -D-galactopyranoside **2** was obtained with a 90% yield following the same protocol, starting from β -methyl D-galactoside. The TMS-protected derivative was then coupled to the fatty carboxylic acid by esterification mediated by 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI) and 4-(dimethylamino)pyridine (DMAP), leading to methyl 6-*O*-acyl-2,3,4-tri-*O*-trimethylsilyl-D-glycopyranosides **3–7** with good to excellent yields (70–88% yield, depending on the alkyl chain and the sugar head group). After quantitative deprotection of the TMS groups using Dowex- H^+ resin and purification on normal phase chromatography, methyl 6-*O*-acyl- α -D-glycopyranosides with C6, C8, C10 and C12 alkyl chains **8–11** and methyl 6-*O*-octanoyl- β -D-galactopyranoside **12** were obtained at the 2g-scale with good to excellent yields (71–83%). The high purity of each intermediate and final compound was confirmed by 1H and ^{13}C NMR (reported in the Supplementary Materials) and HRMS.



Scheme 1. Synthesis of methyl 6-*O*-acyl-*D*-glycopyranosides.


For the sake of brevity, from now on, methyl 6-*O*-acyl- α -*D*-glycopyranosides will be named Me α GlcO(C=O)C_{*n*} (*n* = 5, 7, 9, 11) and Me β GalO(C=O)C₇, respectively, whereas alkyl (methyl *D*-glycopyranosid)uronates will be called Me α Glc(C=O)OC_{*n*} (*n* = 6, 8, 10, 12) and Me β Gal(C=O)OC₈.

2.2. Solubility

Apart from the alkyl chain length, a cornerstone surfactant characteristic, it is generally believed that for sugar-based surfactants in particular, variations in physicochemical properties, including solubility, are often caused by differences in the molecular shape and intra/intermolecular hydrogen bonding [35]. *T_k* represents the temperature above which micelles can form. Below *T_k*, surfactants in hydrated solid form coexist with monomers at their solubility limit, which is visually determined at the transparency limit of the observed solutions. For example, for alkyl glucosides, the *T_k*s of α -anomers are higher than those of the β -anomers and the β -anomers are much more soluble in water than the α -anomers [3]. Table 1 presents the results obtained from both the observation and DSC analysis. In Figure 3, *T_k* is plotted vs. the alkyl chain length for the various studied surfactant families. The results show that, although important, the alkyl chain length is far from the only relevant factor in *T_k* values for sugar-based surfactants. Indeed, the configuration at the anomeric center, as well as the relative orientation of one -OH of the polar head group can also have an important impact.

Table 1. Physical parameters related to the solubility properties (Krafft temperatures, solubility limits) for all studied molecules.

Family	Molecule	Tk (°C)	Solubility Limit (S/mM)	State
Alkyl glycosides	C8 α Glc	39 \pm 1	insoluble at RT; >100 at T > Tk	Powder
	C8 β Glc	<4	>100 at RT	Powder
	C8 α Gal	<4	>100 at RT	Highly viscous liquid
	C8 β Gal	29.0 \pm 0.5	10 < S < 20 at RT; >100 at T > Tk	Powder
Alkyl uronates	Me α Glc(C=O)OC6	<4	\leq 50 at RT	Wax
	Me α Glc(C=O)OC8	<4	15~20 at RT	Wax
	Me α Glc(C=O)OC10	<4	0.5~0.75 at RT	Powder
	Me α Glc(C=O)OC12	22.1 \pm 0.2	0.02~0.05 at RT	Powder
	Me β Gal(C=O)OC8	31.6 \pm 0.9	insoluble at RT; >100 at T > Tk	Powder
Acyl glycosides	Me α GlcO(C=O)C5	<4	<40 at RT	Oil
	Me α GlcO(C=O)C7	<4	\leq 10 at RT	Powder
	Me α GlcO(C=O)C9	26.5 \pm 0.2	insoluble at RT; \leq 1 at T > Tk	Powder
	Me α GlcO(C=O)C11	42.9 \pm 0.1	insoluble at RT; \leq 0.1 at T > Tk	Powder
	Me β GalO(C=O)C7	<4	>100 at RT	Powder

**Figure 3.** Tk as a function of alkyl chain length for studied molecules.

For commercial alkyl glucosides, the Tk decreases in the following sequence: C8 α Glc > C8 β Gal >> C8 β Glc \approx C8 α Gal. All of them possess appreciable solubility limits above their Tks. Many experimental investigations have reported that C8 β Glc is very soluble in water and, therefore, it has a Tk below room temperature, as is also evidenced by our results. The obtained Tk for C8 α Glc is around 40 °C in accordance with previously published values [31,36]. However, as determined by the same experimental protocol, the Tk for C8 β Gal is about 29 °C, much lower than 42 °C derived from the phase diagram reported by Sakya et al. [37] A possible explanation is that Sakya et al. synthesized and purified C8 β Gal by themselves, whereas we carried our measurements on commercially purchased C8 β Gal. Interestingly, in contrast with C8 α Glc, C8 β Gal displays a marked solubility at room temperature even if it is below its Tk. A solution containing 20 mM of C8 β Gal is nearly transparent at room temperature, with only a very low turbidity. For this reason, we assume that many researchers have been able to investigate their aqueous solutions at 25 °C, as mentioned in the introduction. In addition, when cooling the previously heated solutions with the same molecular concentration of 50 mM, we noted that C8 α Glc precipitated much

faster than C8 β Gal. The precipitation of C8 α Glc took place within an hour, whereas no change was observed over one month for the C8 β Gal solution. This may, to some extent, be explained by a more favorable crystal structure for C8 α Glc than C8 β Gal. With regard to C8 α Gal, which is presented in commercial samples as a highly viscous liquid, we expect it to have the weakest crystal structure and a very low Tk value. However, our experimental methods cannot give its exact Tk. Accordingly, one important conclusion can be drawn from our results: for the given type of octyl glycosides, a combination of “axial 4-OH and axial 1-alkyl chain” or “equatorial 4-OH and equatorial 1-alkyl chain”, causes lower Tk values. While a combination like “axial 4-OH and equatorial 1-alkyl chain” or “equatorial 4-OH and axial 1-alkyl chain” can generate higher Tk values. This observation suggests that a different orientation for 4-OH and 1-alkyl chain promotes crystal packing.

After studying commercially available and well-known alkyl glycosides, we focused on the synthesized sugar esters. The observed dissolution phenomena are harder to interpret compared to commercial molecules. This can be a result of more structural contributions such as axial/equatorial 4-OH and 1-OMe, as well as two possible ester orientations. In the case of Me α Glc-derived esters, the solubility decreases, as expected, with the increase of the alkyl chain length. All molecules demonstrated the expected behavior of a lower solubility above their Tk, compared to commercial C8 α Glc and C8 β Glc. Moreover, both octyl glucuronate (Me α Glc(C=O)OC8) and octanoyl glycoside (Me α GlcO(C=O)C7) derivatives showed lower Tk than C8 α Glc. A possible interpretation is that compared to the ether linker, the ester linker creates more disbalance toward hydrogen acceptor groups, which would not find hydrogen donor partners in the crystal structure but would find them in water, increasing aqueous solubility. As highlighted in Table 1, the alkyl (methyl α -D-glycopyranosid)uronates tend to show lower Tk values than their homologous methyl 6-O-acyl- α -D-glycopyranosides. This result is also in accordance with how the molecule presents when its pure form is exposed to some limited atmospheric water. Indeed, we could expect that a molecule showing up as a liquid or paste will likely have a lower Tk compared to a molecule presenting as a crystal in those circumstances. For instance, Me α Glc(C=O)OC8 presents as a semi-solid grease while Me α GlcO(C=O)C7 is a solid powder. No obvious structural interpretation arises regarding this difference. One could speculate that the C=O group in methyl 6-O-acyl- α -D-glycopyranosides is more available to form tight hydrogen bonding networks compared to alkyl (methyl α -D-glycopyranosid)uronates, where it lies closer to the hydrogen donors on the polar head.

For methyl 6-O-acyl- α -D-glycopyranosides, Tk is found to increase with the alkyl chain length, as is the case for alkyl α -glucopyranosides [38]. Moreover, according to Otto et al. [20], the 6-O-octanoyl- β -glucose, above its Tk, has a solubility limit of 25 mg/mL (>80 mM), which is much higher than that of our Me α GlcO(C=O)C7 (<10 mM). This indicates that replacing the anomeric -OH of the glucose with a methyl ether group increases hydrophobicity, making Me α GlcO(C=O)C7 less soluble in water.

In the case of Me β Gal-derived esters, both methyl octyl galacturonate (Me β Gal(C=O)OC8) and methyl 6-O-octanoyl galactoside (Me β GalO(C=O)C7) derivatives have high solubility limits above their Tks, not in line with Me α Glc-derived esters.

The measured Tk for Me β GalO(C=O)C7 is a little higher than that of C8 β Gal. Note that contrary to Me α Glc-derived C7/C8 esters, a carbonyl group closer to the sugar residue is associated with a lower ability to dissolve the molecule at room temperature. One hypothesis is that the polar alcohol group in axial orientation leads to a more compact polar head for Gal surfactants, shielding some polar surface area from contact with the solvent. To have a molecular perspective on this hypothesis, we carried out a comparison with the Me β GlcO(C=O)C8 equivalent. Electrostatic potential maps were obtained with Jmol software after a semi-empirical geometry optimization [39]. The results are displayed in Figure 4.




Figure 4. A 3D representation of (a) Me β GlcO(C=O)C8 and (b) Me β GalO(C=O)C8.

It can be seen that the polar head has a more compact shape; the axial OH is more restricted and has less potential for solvent exposure, supporting a slightly higher hydrophobicity.

2.3. Surface Activity

The aqueous tensiometric properties of all the studied molecules are listed in Table 2. CMCs are determined from the inflections in the plots of equilibrium surface tension against the logarithm of the concentration. For the interested reader, the Gibbs-free energy of micellization, ΔG_{mic} , the Gibbs-free energy of adsorption, ΔG_{ads} , and the efficiency of surfactant adsorption pC_{20} were calculated based on their definitions [40], as presented in Table 2.

Table 2. Surface tension data at $T > T_k$ for different molecules.

Family	Molecule	CMC (mM)	γ_{cmc} (mN/m)	A _{min} (Å ² /molecule)	ΔG_{mic} (kcal/mol)	ΔG_{ads} (kcal/mol)	C ₂₀ (mM)	pC ₂₀	T (°C)
Alkyl glucosides	C8 α Glc	15.5 \pm 0.5	30.7 \pm 0.3	43 \pm 1.8	−4.7	−7.0	2.8	2.6	50
	C8 β Glc	21.2 \pm 0.8	31.0 \pm 0.15	38.5 \pm 1.9	−4.1	−6.4	2.7	2.6	25
	C8 α Gal	51.7 \pm 1.5	29.5 \pm 0.2	45.8 \pm 0.9	−3.6	−6.4	3.8	2.4	25
	C8 β Gal	20 \pm 1	28.5 \pm 0.5	46.3 \pm 1.7	−4.4	−7.1	3.2	2.5	40
Alkyl uronates	Me α Glc(C=O)OC6	55 \pm 5	30.8 \pm 0.8	39.2 \pm 2.9	−3.6	−5.9	5.9	2.2	25
	Me α Glc(C=O)OC8	6 \pm 0.5	29.2 \pm 0.35	41 \pm 1	−4.8	−7.3	0.5	3.3	25
	Me α Glc(C=O)OC10	0.65 \pm 0.5	28.1 \pm 0.1	42.4 \pm 2	−5.9	−8.6	0.05	4.3	25
	Me α Glc(C=O)OC12	0.056 \pm 0.004	28.3 \pm 0.3	45.6 \pm 5	−7.2	−10.1	0.004	5.4	25
	Me β Gal(C=O)OC8	6.9 \pm 0.4	29.4 \pm 0.36	42.9 \pm 3.9	−5.0	−7.4	1.2	2.9	40
Acyl glycosides	Me α GlcO(C=O)C5	105 \pm 5	31 \pm 0.1	40.1 \pm 0.3	−3.3	−5.6	11.2	2.0	25
	Me α GlcO(C=O)C7	9.8 \pm 0.76	29.5 \pm 0.5	39 \pm 2	−4.5	−6.9	1.0	3.0	25
	Me α GlcO(C=O)C9	0.92 \pm 0.08	27.9 \pm 0.05	41.8 \pm 0.9	−5.8	−8.4	0.08	4.1	30
	Me α GlcO(C=O)C11	0.08 \pm 0.005	26.7 \pm 0.25	45.8 \pm 2.8	−7.7	−10.4	0.007	5.1	50
	Me β GalO(C=O)C7	8.5 \pm 0.5	28.3 \pm 0.35	42.7 \pm 2.4	−4.6	−7.3	0.7	3.1	25

Firstly, in the case of octyl D-glycosides, our measured CMC for C8 β Glc, 21.2 mM, was consistent with the published values (18–26 mM), despite the numerous experimental methods used by the different authors [41–45]. However, there were discrepancies in the literature for C8 α Glc, C8 α Gal, and C8 β Gal regarding both CMC and γ_{cmc} . Matsumura et al. [32] measured the static surface tension of C8 α Glc and C8 β Gal by the Wilhelmy plate method at 25 °C. The authors have reported, respectively, CMC values of 12 mM and 16 mM. Both showed a γ_{cmc} of 30.5 mN/m. We measured the CMC values of 15.5 mM and 20 mM, respectively, higher than the published ones, as well as γ_{cmc} 30.7 mN/m and 28.5 mN/m, respectively, indicating a higher effectiveness of surface tension reduction for C8 β Gal. We questioned their results as the measurements were performed below their T_k

(39 °C and 29 °C, respectively, cf. Table 1). Schmidt-Lassen et al. [33] measured the CMC values for C8 β Gal and C8 α Gal by using isothermal titration calorimetry (ITC) and found that CMC values at 25 °C were 31.7 ± 0.7 mM for C8 β Gal and 30.2 ± 0.4 mM for C8 α Gal, showing no evident anomeric effect on the CMC. In contrast, we found CMCs of 20 mM for C8 β Gal and 51.7 mM for C8 α Gal, respectively, indicating that the anomeric orientation has a significant impact on the CMC here. Because the limited aqueous solubility of C8 β Gal was mentioned, again, we question the results published in their study.

After comparing our measurements with published ones, we focused on the structural interpretations arising from our own measurements. The order of their CMC for the 4 octyl glycosides was: C8 α Gal > C8 β Glc ~ C8 β Gal > C8 α Glc. Here, even at a constant alkyl chain length, a lower CMC correlates with a higher T_k . Apart from the alkyl chain length, the most obvious factor, it is known that polar head hydrophilicity can also significantly influence CMC, typically increasing it for more hydrophilic polar heads [6]. With all else equal, a higher CMC leads to a decrease in T_k due to the thermodynamic relationship between the two. Thus, the observed T_k could also reflect polar head hydrophilicity. As for γ_{cmc} , it was found that octyl D-galactosides can more effectively reduce the surface tension of pure water than octyl glucosides. Our A_{min} measurements suggest that the anomeric effect is more evident for C8 α - and C8 β Glc than for C8 α - and C8 β Gal. In addition, the latter shows a greater A_{min} than the former, in agreement with Razafindralambo et al. [14]. It was said that an axial -OH at the 4-position (Gal) gives a larger minimum area occupied per molecule at the water surface. However, as the A_{min} depends strongly on the way of plotting the γ vs. $\log C$ curves, different authors have demonstrated different A_{min} values. Matsumura et al. [32] reported a range of 47–54 Å²/molecule for C8–C12 alkyl D-glucosides and galactosides. Their study did not show any great difference between their α - and β -anomers but a slightly larger A_{min} for C8- and C10 β Gal than for the C8- and C10 α Glc. Kjellin et al. [23] showed an A_{min} of 38 Å²/molecule for C8 β Glc whereas an A_{min} of 42 Å²/molecule was given for C8 β Glc by Shinoda et al. [46]. Therefore, our results seem to be in good agreement with the literature.

The influence of the head saccharidic group of our new surfactants on the surface activity can also be analyzed. Figure 5 (top) represents the surface tension vs. \log concentration plots for both families of Me α Glc-derived esters. Firstly, whatever the orientation of the ester group (whatever the family studied), the esters of glucose can efficiently reduce the surface tension of pure water with the obtained low γ_{cmc} values of 26–32 mN/m. It seems that the ester group orientation does not impact γ_{cmc} much. In general, it can be seen that γ_{cmc} slightly decreases with the alkyl chain length. Those results are in line with the identified trends [6] where γ_{cmc} appears to decrease asymptotically with the alkyl chain length. The suggested explanation is that as the alkyl chain length increases, alkyl groups get closer and closer to completely covering the surface, making it exhibit an alkane-like γ_{cmc} . The only exception comes with Me α Glc(C=O)OC12, which is very close to that of Me α Glc(C=O)OC10, very likely because it is challenging to measure a true γ_{cmc} for Me α Glc(C=O)OC12 as it precipitates easily. A similar phenomenon was observed by Otto et al. [20], who reported a higher γ_{cmc} for 6-O-decanoyl- β -D-glucopyranose (C10) than for 6-O-octanoyl- β -D-glucopyranose (C8).

A common conclusion in the literature is that the alkyl chain length is not an important factor in determining the A_{min} of n-alkyl glucosides at the air–water interface [3]. This observation can probably be extrapolated to sugar esters. Nevertheless, even though A_{min} is mainly decided by the hydrophilic group [47], the alkyl chain flexibility can contribute to whether molecules loosely or compactly pack at the interface [48]. It is reasonable to assume that given the higher number of rotational degrees of freedom, a longer saturated alkyl chain is more flexible, and based on this argument, we hypothesize that surfactants with longer alkyl chains at a constant polar head can exhibit the A_{min} increase. Our measurements are consistent with such a hypothesis, with a slight increase of A_{min} with alkyl chain lengths for both families. A similar trend was mentioned in the literature for other surfactant families, such as bolaform quaternary ammonium surfactants for

which the A_{min} increases with the increase in the spacer carbon number ($n < 10$) [49]. Nevertheless, it must be noted that for some polyoxyethylene alkyl ethers, A_{min} decreases with an increasing carbon number but increases with the oxyethylene unit length [47,50,51]. Given that contrary to rigid polar rings of sugar-based surfactants, the polar heads of polyoxyethylene alkyl ethers are themselves flexible, their conformation may change with longer alkyl chains. Thus, the above analysis may not apply in their case.




Figure 5. Surface tension as a function of concentration for methyl alkyl α -D-glucuronates at 25 °C (empty markers, $\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_n$, with n being an even number) and methyl 6-O-acyl- α -D-glucosides (filled markers, $\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_n$, with n being an odd number) at 25 °C (C5 and C7), 30 °C (C9) and 50 °C (C11) (top) and the variation of the log CMC with the carbon numbers of the alkyl chain for alkyl methyl α -D-glucuronates ($\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_n$, Δ) and methyl 6-O-acyl- α -D-glucosides ($\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_n$, \square) (bottom). CMC is expressed in mol/L.

When it comes to CMC, the alkyl chain length, as well-known for other surfactants [6], is the most relevant structural parameter. The dependence of the log CMC on the number of carbon atoms in the alkyl chain for two families, generally, obeys the expected empirical equation [21]:

$$\text{Log CMC} = A - Bn \quad (1)$$

where n represents the carbon numbers (the length) in the alkyl chain; A and B are empirical constants, which are, respectively, related to the contributions of the hydrophilic head group and a single hydrophobic methylene group to the change in free energy for micellization. Plots of log CMC values as functions of the alkyl chain length are depicted in Figure 5 (bottom).

The results show that a linear relationship between the alkyl chain length and the log CMC exists for both ester families. For alkyl D-glucuronates derivatives ($\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_n$), the values of A and B calculated are 0.5 and 1.7, while for acyl D-glucosides derivatives ($\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_n$), A is 0.5 and B is 1.5. The slightly higher B value in the case of alkyl D-glucuronates suggests a more rapid decrease in CMC when one carbon is added. It is important to note that the $\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_n$ series shows even-numbered carbon atoms while the $\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_n$ series has odd-numbered carbon atoms in the noted C_n part. The amphiphilic molecules can be, in a simplistic way, depicted as the combination of a polar head group with a lipophilic tail. In the case of both families, the ester linkage is attributed to the polar moiety, even if it is not a part of the sugar: for acyl glycosides, this functional group is not located on a carbon atom of the sugar, whereas for the alkyl D-glucuronate, the saccharidic C6 is oxidized, leading to a difference in the structure (and particularly in the number of carbon atoms) of the polar head. We assume that this slight difference has a negligible effect on the hydrophilicity of the head group. On the other hand, the alkyl D-glucuronates series ($\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_n$) exhibits a slightly lower CMC and γ_{cmc} compared to the acyl D-glucosides series ($\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_n$), which can be partially associated with the fact that the former always has one more carbon in its lipophilic moiety than the latter.

Some published CMC values are also worth comparing to the measurements from the present work. Otto et al. [20] obtained a CMC of 18 mM for 6-*O*-octanoyl-D-glucopyranose, noticeably higher than the CMC of $\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_7$ (9.8 mM). In addition, Blecker et al. [17] also reported a CMC of 10.68 mM for octyl D-glucuronate compared to the 6 mM value we observed for the methyl octyl α -D-glucuronate ($\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_8$). Both comparisons indicate that the nonpolar axial methyl group at the 1-position brings additional hydrophobicity to the polar head.

For the first time, Razafindralambo et al. [52] studied the effect of the ester orientation (Su-O-CO-R or Su-CO-O-R) on the surface properties of similar molecules: D-glucosyl octanoate and octyl D-glucuronate. Their analysis was that octyl D-glucuronate exhibits a stronger hydrophobic character because its carbonyl group was closer to the head group, making the head more polar and resulting in stronger intermolecular interactions between the heads. The CMC values at 25 °C for octyl D-glucuronate and D-glucosyl octanoate were reported to be 10.7 and 19.1 mM, respectively, which were almost twice as large as the CMC values for $\text{Me}\alpha\text{Glc}(\text{C}=\text{O})\text{OC}_8$ and $\text{Me}\alpha\text{GlcO}(\text{C}=\text{O})\text{C}_7$. It is worth noting that the Razafindralambo molecules are a mixture of both α and β anomers, with the β -D-glucose derivative being more hydrophilic. In addition, they obtained a larger A_{min} for octyl D-glucuronate, which was supported by a computational approach. Based on those results, the authors suggested that a shorter distance between the carbonyl group and the sugar head introduces a tilt between the polar head and the alkyl chain, resulting in looser packing at the water surface. For the synthesized surfactants of this work, however, our measurements do not evidence a significant influence of the ester orientation on A_{min} . It is possible that the anomeric methyl group dilutes the influence of the mechanism proposed by Razafindralambo et al. Nevertheless, their proposed mechanism itself remains questionable as the surface is a dynamic environment where molecules not only take different conformations but also different orientations. By comparing our results with

other literature work, we note that the location of the ester group at the head group also influenced the CMC: sugar esters with the carbonyl group located at the 3-position appear more hydrophobic than those with the carbonyl group located at the 6-position. For example, Gou eth et al. [19] reported that 3-*O*-octanoyl-*D*-glucopyranose, which is a mixture of α and β anomers, had a CMC of 0.61 mM while Savellie et al. [21] reported a CMC of 1.6 mM. Both studies evidenced a lower CMC than that for 6-*O*-octanoyl- β -*D*-glucopyranose and our octyl derivative esters, as discussed above. Moreover, these molecules (Me α Glc-(C=O)OC8 and Me α GlcO(C=O)C7) show much lower CMC values compared to the commercial C8 α Glc and C8 β Glc, as shown in Figure 6. The results indicate that both the methyl group at the 1-position and the transformation of 6-OH into the ester group increase the hydrophobicity but do not influence γ_{cmc} significantly.




Figure 6. Surface tension as a function of the concentration for α -octyl glucoside (C8 α Glc, \circ) at 50 $^{\circ}$ C, β -octyl glucoside (C8 β Glc, \bullet) at 25 $^{\circ}$ C, octyl α -methyl glucuronate (Me α Glc(C=O)OC8, \diamond) at 25 $^{\circ}$ C and 6-octanoyl- α -methyl glucoside (Me α GlcO(C=O)C7, \blacklozenge) at 25 $^{\circ}$ C.

Our A_{min} measurements (see Table 2) indicate a more compact packing for the synthesized esters at the interface of the air–aqueous solution. This might be related to the stronger intermolecular interactions between their sugar heads due to the presence of the ester group.

In the case of Me β Gal-based esters derivatives (Figure 7), the octyl *D*-galacturonate derivative (Me β Gal(C=O)OC8) also showed a lower CMC than its homologous octanoyl *D*-galactoside derivative (Me β GalO(C=O)C7), which is consistent with Me α Glc-derived C7/C8 esters. Given that the actual alkyl chain of Me β Gal(C=O)OC8 contains 8 carbons vs. 7 for Me β GalO(C=O)C7, and that the alkyl chain length is the major parameter relevant to CMC, the observed CMC difference is in line with expectations. It can also be noticed that the CMCs observed for the esters (6–9.8 mM) are lower than those of corresponding octyl *D*-galactosides (20–51.7 mM). Given the higher observed A_{min} for octyl *D*-galactosides, the shape of the esters' polar heads may allow micellization more readily.




Figure 7. Surface tension as a function of concentration for β -octyl galactoside ($C8\beta Gal$, \circ) at $40\text{ }^{\circ}C$, α -octyl galactoside ($C8\alpha Gal$, \bullet) at $25\text{ }^{\circ}C$, octyl β -methyl galacturonate ($Me\beta Gal(C=O)OC8$, \diamond) at $40\text{ }^{\circ}C$ and 6-octanoyl- β -methyl galactoside ($Me\beta GalO(C=O)C7$, \blacklozenge) at $25\text{ }^{\circ}C$.

Overall, the main conclusions that arise by comparing the solubility and surface activity for all the synthesized C7/C8 esters are listed below:

- $Me\beta Gal$ -based esters show a much larger solubility limit than $Me\alpha Glc$ -based esters, suggesting that the solubility strongly depends on the head group configuration;
- Ester group orientation also influences the dissolution of $Me\beta Gal$ -based esters and $Me\alpha Glc$ -based esters;
- CMC is influenced by the ester orientation following the relationship: $Me\alpha Glc(C=O)OC8 < Me\beta Gal(C=O)OC8 < Me\beta GalO(C=O)C7 < Me\alpha GlcO(C=O)C7$. The result suggests that the uronates derivatives exhibit lower CMC values than the acyl D-glycosides derivatives. The effect of ester group orientation on CMC was more evident for $Me\alpha Glc$ -derived esters than for $Me\beta Gal$ -derived esters;
- γ_{cmc} is very similar for all these esters, suggesting that the ester linkage orientation does not play an important role in the effectiveness of reducing surface tension.

3. Materials and Methods

3.1. General

Octyl α -D-glucopyranoside (>99%) and octyl β -D-glucopyranoside (>99%) were both purchased from Anatrace (Maumee, OH, USA). Octyl α -D-galactopyranoside and octyl β -D-galactopyranoside were purchased from Carbosynth (Compton, UK). They were all used without further purification. The deionized water with a resistivity of 18.2 M Ω cm was produced by a lab purification chain provided by Aquadem/Veolia Water STI (Saint-Maurice, France) and used to prepare all the solutions.

Alkyl (methyl D-glycopyranosid)uronates were synthesized using the two-step procedure developed in our lab, as previously described [29]. Briefly, the primary hydroxyl group of free methyl D-glycopyranosides was selectively and quantitatively oxidized by using 2,2,6,6-tetramethyl-1-piperidinyloxy free radical (TEMPO). Hydrophobic chains of different lengths were then introduced by acid-mediated esterification with fatty alcohols

(hexyl to dodecyl alcohols) leading to the desired alkyl uronates with moderate to good yields (49–63%).

3.2. Synthesis of Methyl 6-O-acyl-D-glycopyranosides

3.2.1. General

All chemicals were purchased from Fisher Scientific (Illkirch, France), and Merck Sigma Aldrich (St Quentin Fallavier, France), and used as received. Mass analyses were performed on a Waters spectrometer (SINAPT TG2SI, Manchester, UK) using electrospray ionization (Z-Spray). The NMR analysis was performed on a BRUKER NMR spectrometer (Bruker, Wissembourg, France) operating at 400 MHz for ^1H and 100 MHz for ^{13}C . Samples of silylated products were prepared in deuterated chloroform (CDCl_3) while samples of free esters were dissolved in $\text{DMSO}-d_6$.

3.2.2. General Procedure for One-Pot Protection/Selective Deprotection

N,O-bis(trimethylsilyl)acetamide (28.7 g, 4.3 eq.) and tetrabutylammonium fluoride (235 mg, 0.03 eq.) were added to a solution of methyl α -D-glucopyranoside (9.30 g, 32.9 mmol) in dry pyridine (7.5 mL). After 1.5 h of stirring at room temperature, the reaction was quenched by adding isopropanol (6.25 mL) and methanol (450 mL). After cooling to 0 °C, K_2CO_3 (2.28 g, 0.5 eq.) was added, the mixture was stirred for 30 min at 0 °C, and then neutralized with 2.5 mL of acetic acid. After concentration, the crude was extracted with Et_2O , washed with water, dried over Na_2SO_4 , and concentrated to give methyl 2,3,4-tri-*O*-trimethylsilyl- α -D-glucopyranoside **1** as a white powder (10.94 g, yield 80%). The same procedure was applied to methyl β -D-galactopyranoside affording methyl 2,3,4-tri-*O*-trimethylsilyl- β -D-galactopyranoside **2** as a white powder (12.13 g, yield 90%).

3.2.3. General Procedure for Esterification

To a solution of methyl 2,3,4-tri-*O*-trimethylsilyl- α -D-glycopyranoside **1** or **2** (7.00 g, 17.1 mmol) in anhydrous dichloromethane (80 mL), 4-dimethylaminopyridine (1.04 g, 0.5 eq.), desired fatty acid (37.5 mmol, 2.2 eq.), and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (5.29 g, 2 eq.) were successively added and the mixture was stirred at room temperature for 6 h. After concentration, the crude was purified by normal phase column chromatography on silica gel (mobile phase: cyclohexane/ EtOAc 9:1) to give corresponding silylated sugar esters **3–7**.

3.2.4. General Procedure for Deprotection

To a solution of the as-obtained silylated ester glycoside, **3–7** (6 g) in $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 1:1 (100 mL), 3 g of Dowex- H^+ resin was added. The suspension was stirred for 30 min at room temperature, then the resin was eliminated by filtration, and the filtrate was concentrated. The residue was purified by column chromatography on silica gel (mobile phase: EtOAc/MeOH 98:2) to give pure corresponding methyl 6-*O*-acyl D-glycosides **8–12**.

3.3. Compound Characterization Data

Methyl 2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 1. White powder. Yield 80% from methyl α -D-glucopyranoside following the general procedure. ^1H and ^{13}C NMR spectra in accordance with the literature [53].

Methyl 2,3,4-tri-O-trimethylsilyl- β -D-galactopyranoside 2. White powder. Yield 90% from methyl β -D-galactopyranoside following the general procedure. ^1H and ^{13}C NMR spectra in accordance with the literature [54].

Methyl 6-O-hexanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 3. Colorless syrup. Yield 72% from the methyl 2,3,4-tri-*O*-trimethylsilyl- α -D-glucopyranoside **1** following the general procedure. ^1H NMR (400 MHz, CDCl_3), δ 0.11–0.16 (m, 27H, $(\text{CH}_3)_3\text{-Si}$), 0.90 (t, $J_{\text{CH}_3,\text{CH}_2} = 6.9$ Hz, 3H, $(\text{CH}_2)\text{-CH}_3$), 1.30–1.35 (m, 4H, $\text{CH}_2\text{-}(\text{CH}_2)_2\text{-CH}_3$), 1.60–1.68 (m, 2H, $\text{CO-CH}_2\text{-CH}_2$), 2.35 (o, $J = 7.7$ Hz, 2H, $\text{CO-CH}_2\text{-CH}_2$), 3.35 (s, 3H, O-CH_3), 3.42–3.49 (m, 2H, H-2,H-5), 3.68–3.76 (m, 2H, H-3, H-4), 4.03 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.9$ Hz, $J_{\text{H-5,H-6}'} = 5.4$ Hz

H-6'), 4.40 (dd, 1H, $J_{H-6,H-6'} = 11.9$ Hz, $J_{H-5,H-6} = 2.2$ Hz, H-6), 4.70 (d, 1H, $J_{H-1,H-2} = 3.7$ Hz, H-1). ^{13}C NMR (100 MHz, CDCl_3) δ 0.5–1.5 (3s, $(\text{CH}_3)_3\text{-Si}$), 13.9 ($\text{CH}_2\text{-CH}_3$), 22.3–31.3 (2s, $\text{CH}_2\text{-(CH}_2)_2\text{-CH}_3$), 24.5 ($\text{CO-CH}_2\text{-CH}_2\text{-(CH}_2)_2$), 34.2 ($\text{CO-CH}_2\text{-CH}_2$), 54.8 (O-CH_3), 63.3 (C-6), 69.4 (C-4), 72.5 (C-2), 73.7 (C-5), 75.1 (C-3), 99.8 (C-1), 173.6 (CO).

Methyl 6-O-octanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 4. Colorless syrup. Yield 77% from the methyl 2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside **1** following the general procedure. ^1H NMR (400 MHz, CDCl_3) δ 0.10–0.15 (3 s, 27H, $(\text{CH}_3)_3\text{-Si}$), 0.85 (t, $J_{\text{CH}_3,\text{CH}_2} = 6.9$ Hz, 3H, $\text{CH}_2\text{-CH}_2\text{-CH}_3$), 1.20–1.31 (m, 8H, $\text{CH}_2\text{-(CH}_2)_4\text{-CH}_3$), 1.58–1.63 (m, 2H, $\text{CO-CH}_2\text{-CH}_2\text{-CH}_2$), 2.32 (m, 2H, $\text{CO-CH}_2\text{-CH}_2$), 3.32 (s, 3H, O-CH_3), 3.42–3.48 (m, 2H, H-2, H-5), 3.67–3.75 (m, 2H, H-3, H-4), 4.02 (dd, 1H, $J_{H-6,H-6'} = 11.9$ Hz, $J_{H-5,H-6'} = 5.4$ Hz H-6'), 4.37 (d, 1H, $J_{H-6,H-6'} = 11.8$ Hz), 4.62 (d, 1H, $J_{H-1,H-2} = 3.7$ Hz, H-1). ^{13}C NMR (100 MHz, CDCl_3) δ 0–1.5 (3s, $(\text{CH}_3)_3\text{-Si}$), 14.0 ($\text{-CH}_2\text{-CH}_3$), 22.5–31.5 (4s, $\text{CH}_2\text{-(CH}_2)_4\text{-CH}_3$), 24.8 ($\text{CO-CH}_2\text{-CH}_2\text{-CH}_2$), 34.2 ($\text{CO-CH}_2\text{-CH}_2$), 54.7 (O-CH_3), 63.3 (C-6), 69.5 (C-4), 72.5 (C-2), 73.7 (C-5), 75.1 (C-3), 99.7 (C-1), 173.5 (CO).

Methyl 6-O-decanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 5. Colorless syrup. Yield 88% from the methyl 2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside **1** following the general procedure. ^1H NMR (400 MHz, CDCl_3) δ 0.11–0.16 (3 s, 27H, $(\text{CH}_3)_3\text{-Si}$), 0.85 (t, $J_{\text{CH}_3,\text{CH}_2} = 6.9$ Hz, 3H, $\text{CH}_2\text{-CH}_3$), 1.19–1.35 (m, 12H, $\text{CH}_2\text{-(CH}_2)_6\text{-CH}_3$), 1.55–1.65 (m, 2H, $\text{CO-CH}_2\text{-CH}_2$), 2.37 (td, 2H, $\text{CO-CH}_2\text{-CH}_2$), 3.35 (s, 3H, O-CH_3), 3.43–3.5 (m, 2H, H-2, H-5), 3.68–3.76 (m, 2H, H-3, H-4), 4.03 (dd, 1H, $J_{H-6,H-6'} = 11.8$ Hz, $J_{H-5,H-6'} = 5.3$ Hz, H-6'), 4.37 (dd, 1H, $J_{H-6,H-6'} = 11.8$ Hz, $J_{H-5,H-6} = 2.3$ Hz H-6), 4.60 (d, 1H, $J_{H-1,H-2} = 3.7$ Hz, H-1). ^{13}C NMR (100 MHz, CDCl_3) δ 0–1.5 (3s, $(\text{CH}_3)_3\text{-Si}$), 14.0 ($\text{-CH}_2\text{-CH}_3$), 22.5–32 (6s, $\text{CH}_2\text{-(CH}_2)_6\text{-CH}_3$), 24.7 ($\text{CO-CH}_2\text{-CH}_2\text{-CH}_2$), 34.2 ($\text{CO-CH}_2\text{-CH}_2$), 55.0 (O-CH_3), 63.2 (C-6), 69.4 (C-4), 72.5 (C-2), 73.8 (C-5), 75.0 (C-3), 99.8 (C-1), 173.5 (CO).

Methyl 6-O-dodecanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 6. Colorless syrup. Yield 70% from the methyl 2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside **1** following the general procedure. ^1H NMR (400 MHz, CDCl_3) δ 0.14–0.19 (3s, 27H, $(\text{CH}_3)_3\text{-Si}$), 0.75 (t, $J_{\text{CH}_3,\text{CH}_2} = 6.9$ Hz, 3H, $\text{CH}_2\text{-CH}_2\text{-CH}_3$), 1.20–1.30 (m, 16H, $\text{CH}_2\text{-(CH}_2)_8\text{-CH}_3$), 1.61–1.64 (m, 2H, $\text{CO-CH}_2\text{-CH}_2\text{-CH}_2$), 2.35 (td, 2H, CO-CH_2), 3.35 (s, 3H, O-CH_3), 3.43–3.49 (m, 2H, H-2, H-5), 3.68–3.78 (m, 2H, H-3, H-4), 4.05 (dd, 1H, $J_{H-6,H-6'} = 11.8$ Hz, $J_{H-5,H-6'} = 5.4$ Hz, H-6'), 4.39 (dd, 1H, $J_{H-6,H-6'} = 11.8$ Hz, $J_{H-5,H-6} = 2.2$ Hz, H-6), 4.62 (d, 1H, $J_{H-1,H-2} = 3.6$ Hz, H-1). ^{13}C NMR (100 MHz, CDCl_3) δ 0–1.5 (3s, $(\text{CH}_3)_3\text{-Si}$), 14.0 ($\text{-CH}_2\text{-CH}_3$), 22.5–21.5 (8s, $\text{CH}_2\text{-(CH}_2)_8\text{-CH}_3$), 24.5 ($\text{CO-CH}_2\text{-CH}_2$), 34.3 (CO-CH_2), 55.0 (O-CH_3), 63.2 (C-6), 69.5 (C-4), 72.6 (C-2), 73.8 (C-5), 75.0 (C-3), 99.8 (C-1), 173.4 (CO).

Methyl 6-O-octanoyl-2,3,4-tri-O-trimethylsilyl- β -D-galactopyranoside 7. Colorless syrup. Yield 83% from the methyl 2,3,4-tri-O-trimethylsilyl- β -D-galactopyranoside **2** following the general procedure. ^1H NMR (400 MHz, CDCl_3) δ 0.12 (3s, 27H $(\text{CH}_3)_3\text{-Si}$), 0.85 (t, $J_{\text{CH}_3,\text{CH}_2} = 6.9$ Hz, 3H, $\text{CH}_2\text{-CH}_2\text{-CH}_3$), 1.20–1.31 (m, 8H, $\text{CH}_2\text{-(CH}_2)_4\text{-CH}_3$), 1.61 (p, 2H, $J = 7.4$ Hz $\text{CO-CH}_2\text{-CH}_2\text{-CH}_2$), 2.30 (t, 2H, $J = 7.5$ Hz, $\text{CO-CH}_2\text{-CH}_2$), 3.40 (dd, 1H, $J_{H-2,H-3} = 9.3$ Hz, $J_{H-3,H-4} = 2.7$ Hz, H-3), 3.47 (s, 3H, O-CH_3), 3.64 (m, 2H, H-2, H-5), 3.74 (d, 1H, $J_{H-3,H-4} = 2.3$ Hz, H-4), 4.06 (d, 1H, $J_{H-1,H-2} = 7.5$ Hz, H-1), 4.13 (dd, 1H, $J_{H-6,H-6'} = 11.0$ Hz, $J_{H-5,H-6} = 6.5$ Hz, H-6), 4.23 (dd, 1H, $J_{H-6,H-6'} = 11.0$ Hz, $J_{H-5,H-6'} = 6.6$ Hz, H-6'), ^{13}C NMR (100 MHz, CDCl_3) δ 0.6, 0.7, 0.8 (3s, $(\text{CH}_3)_3\text{-Si}$), 14.2 ($\text{-CH}_2\text{-CH}_3$), 22.7, 29.0, 29.2, 31.7 (4s, $\text{CH}_2\text{-(CH}_2)_4\text{-CH}_3$), 25.1 ($\text{CO-CH}_2\text{-CH}_2\text{-CH}_2$), 34.4 ($\text{CO-CH}_2\text{-CH}_2$), 57.2 (O-CH_3), 63.0 (C-6), 71.9 (C-4), 72.0 (C-2), 72.2 (C-5), 75.1 (C-3), 105.0 (C-1), 173.6 (CO).

Methyl 6-O-hexanoyl- α -D-glucopyranoside 8. Colorless syrup. Yield 77% from the methyl 6-O-hexanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside **3** following the general procedure for deprotection. ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 0.84 (t, 3H, $J_{\text{CH}_3,\text{CH}_2} = 6.9$ Hz, $\text{CH}_2\text{-CH}_3$), 1.21–1.29 (m, 4H, $\text{-(CH}_2)_2\text{-CH}_3$), 1.44–1.57 (m, 2H, $\text{CO-CH}_2\text{-CH}_2$), 2.28 (t, 2H, $J_{\text{CH}_2,\text{CH}_2} = 7.3$ Hz, CO-CH_2), 3.02–3.09 (m, 1H, H-4), 3.16–3.23 (m, 1H, H-2), 3.25 (s, 3H, O-CH_3), 3.35–3.41 (m, 1H, H-3), 3.48–3.53 (m, 1H, H-5), 4.01 (dd, 1H, $J_{H-6,H-6'} = 11.6$ Hz, $J_{H-5,H-6'} = 6.5$ Hz, H-6'), 4.29 (dd, 1H, $J_{H-6,H-6'} = 11.6$ Hz, $J_{H-5,H-6} = 2.1$ Hz, H-6), 4.52 (d, 1H, $J_{H-1,H-2} = 3.7$ Hz, H-1), 4.82 (d, 1H, $J_{\text{OH},\text{H-}2'} = 6.4$ Hz, OH-2), 4.90 (d, 1H, $J_{\text{OH},\text{H-}3'} = 5.0$ Hz, OH-3), 5.15 (d, 1H, $J_{\text{OH},\text{H-}4'} = 5.8$ Hz, OH-4). ^{13}C (100 MHz, $\text{DMSO-}d_6$) δ 13.9 ($\text{-CH}_2\text{-CH}_3$),

22.9, 24.3, 30.7 (3 s, $-(\text{CH}_2)_3\text{-CH}_3$), 33.6 (CO-CH_2 -), 54.5 (O-CH_3), 63.6 (C-6), 69.7 (C-5), 70.5 (C-4), 71.9 (C-2), 73.3 (C-3), 99.8 (C-1), 173.0 (CO). HRMS (ESI) m/z for $\text{C}_{13}\text{H}_{24}\text{O}_7\text{Na}^+$ $[\text{M} + \text{Na}]^+$ calcd 315.1420, found 315.1424.

Methyl 6-O-octanoyl- α -D-glucopyranoside 9. White powder. Yield 77% from the methyl 6-O-octanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 4 following the general procedure for deprotection. ^1H NMR (400 MHz, DMSO- d_6) δ 0.84 (t, 3H, $J_{\text{CH}_3,\text{CH}_2} = 7.3$ Hz, $\text{CH}_2\text{-CH}_3$), 1.17–1.30 (m, 8H, $-(\text{CH}_2)_4\text{-CH}_3$), 1.47–1.55 (m, 2H, $\text{CO-CH}_2\text{-CH}_2$), 2.28 (t, 2H, $J_{\text{CH}_2,\text{CH}_2} = 7.3$ Hz, CO-CH_2), 3.02–3.09 (m, 1H, H-4) 3.16–3.23 (m, 1H, H-2) 3.25 (s, 3H, O-CH_3), 3.35–3.41 (m, 1H, H-3), 3.50–3.56 (m, 1H, H-5) 4.01 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.7$ Hz, $J_{\text{H-5,H-6}'} = 6.7$ Hz, H-6'), 4.28 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.7$ Hz, $J_{\text{H-5,H-6}} = 1.9$ Hz, H-6), 4.52 (d, 1H, $J_{\text{H-1,H-2}} = 3.6$ Hz, H-1) 4.82 (d, 1H, $J_{\text{OH,H-2}'} = 6.4$ Hz, OH-2), 4.90 (d, 1H, $J_{\text{OH,H-3}'} = 4.9$ Hz, OH-3), 5.15 (d, 1H, $J_{\text{OH,H-4}'} = 5.8$ Hz, OH-4). ^{13}C (100 MHz, DMSO- d_6) δ 14.1 ($-\text{CH}_2\text{-CH}_3$), 22.2, 24.6, 28.5, 28.5, 31.2 (5 s, $(\text{CH}_2)_5\text{-CH}_3$), 33.6 (CO-CH_2 -), 54.4 (O-CH_3), 63.6 (C-6), 69.7 (C-5), 70.5 (C-4), 71.9 (C-2), 73.3 (C-3), 99.8 (C-1), 173.0 (CO). HRMS (ESI) m/z for $\text{C}_{15}\text{H}_{28}\text{O}_7\text{Na}^+$ $[\text{M} + \text{Na}]^+$ calcd 343.1733, found 343.1726.

Methyl 6-O-decanoyl- α -D-glucopyranoside 10. White powder. Yield 81% from the methyl 6-O-decanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 5 following the general procedure for deprotection. ^1H NMR (400 MHz, CD_3OD) δ 0.89 (t, 3H, $J_{\text{CH}_3,\text{CH}_2} = 7.3$ Hz, $\text{CH}_2\text{-CH}_3$), 1.24–1.39 (m, 12H, $-(\text{CH}_2)_6\text{-CH}_3$), 1.62 (p, 2H, $J_{\text{CH}_2,\text{CH}_2} = 7.2$ Hz, $\text{CO-CH}_2\text{-CH}_2$), 2.35 (t, 2H, $J_{\text{CH}_2,\text{CH}_2} = 7.4$ Hz, $\text{CO-CH}_2\text{-CH}_2$), 3.27 (dd, 1H, $J_{\text{H-3,H-4}} = 9.0$ Hz, $J_{\text{H-4,H-5}} = 9.9$ Hz, H-4) 3.37–3.42 (m, 4H, H-2, O-CH_3), 3.61 (t, 1H, $J_{\text{H-2,H-3}} = J_{\text{H-3,H-4}} = 9.3$ Hz, H-3), 3.70 (ddd, 1H, $J_{\text{H-4,H-5}} = 9.9$ Hz, $J_{\text{H-5,H-6}} = 2.0$ Hz, $J_{\text{H-5,H-6}'} = 6.0$ Hz, H-5), 4.19 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.8$ Hz, $J_{\text{H-5,H-6}'} = 6.0$ Hz, H-6'), 4.38 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.8$ Hz, $J_{\text{H-5,H-6}} = 2.1$ Hz, H-6), 4.65 (d, 1H, $J_{\text{H-1,H-2}} = 3.7$ Hz, H-1). ^{13}C (100 MHz, CD_3OD) δ 14.5 ($-\text{CH}_2\text{-CH}_3$), 23.7, 26.1, 30.2, 30.4, 30.4, 30.6, 33.0 (7 s, $-(\text{CH}_2)_7\text{-CH}_3$), 35.0 (CO-CH_2 -), 55.6 (O-CH_3), 64.7 (C-6), 71.0 (C-5), 71.9 (C-4), 73.4 (C-2), 75.0 (C-3), 101.3 (C-1), 175.4 (CO). HRMS (ESI) m/z for $\text{C}_{17}\text{H}_{32}\text{O}_7\text{Na}^+$ $[\text{M} + \text{Na}]^+$ calcd 371.2046, found 371.2046.

Methyl 6-O-dodecanoyl- α -D-glucopyranoside 11. White powder. Yield 71% from the methyl 6-O-dodecanoyl-2,3,4-tri-O-trimethylsilyl- α -D-glucopyranoside 6 following the general procedure for deprotection. ^1H NMR (400 MHz, CD_3OD) δ 0.89 (t, 3H, $J_{\text{CH}_3,\text{CH}_2} = 7.3$ Hz, $\text{CH}_2\text{-CH}_3$), 1.24–1.40 (m, 16H, $-(\text{CH}_2)_8\text{-CH}_3$), 1.62 (p, 2H, $J_{\text{CH}_2,\text{CH}_2} = 7.2$ Hz, $-\text{CO-CH}_2\text{-CH}_2$), 2.35 (t, 2H, $J_{\text{CH}_2,\text{CH}_2} = 7.4$ Hz, CO-CH_2), 3.27 (dd, 1H, $J_{\text{H-3,H-4}} = 9.0$ Hz, $J_{\text{H-4,H-5}} = 10.0$ Hz, H-4) 3.37–3.42 (m, 4H, H-2, O-CH_3), 3.61 (t, 1H, $J_{\text{H-2,H-3}} = J_{\text{H-3,H-4}} = 9.3$ Hz, H-3), 3.70 (ddd, 1H, $J_{\text{H-4,H-5}} = 9.9$ Hz, $J_{\text{H-5,H-6}} = 2.0$ Hz, $J_{\text{H-5,H-6}'} = 6.0$ Hz, H-5), 4.19 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.8$ Hz, $J_{\text{H-5,H-6}'} = 6.0$ Hz, H-6'), 4.38 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.8$ Hz, $J_{\text{H-5,H-6}} = 2.1$ Hz, H-6), 4.65 (d, 1H, $J_{\text{H-1,H-2}} = 3.7$ Hz, H-1). ^{13}C (100 MHz, CD_3OD) δ 14.5 ($-\text{CH}_2\text{-CH}_3$), 23.8, 26.1, 30.2, 30.4, 30.5, 30.6, 30.8, 30.8, 33.1 (9 s, $-(\text{CH}_2)_9\text{-CH}_3$), 35.0 (CO-CH_2 -), 55.6 (O-CH_3), 64.7 (C-6), 71.1 (C-5), 71.9 (C-4), 73.5 (C-2), 75.0 (C-3), 101.3 (C-1), 175.4 (CO). HRMS (ESI) m/z for $\text{C}_{19}\text{H}_{36}\text{O}_7\text{Na}^+$ $[\text{M} + \text{Na}]^+$ calcd 399.2359, found 399.2361.

Methyl 6-O-octanoyl- β -D-galactopyranoside 12. White powder. Yield 83% from the methyl 6-O-octanoyl-2,3,4-tri-O-trimethylsilyl- β -D-galactopyranoside 7 following the general procedure for deprotection. ^1H NMR (400 MHz, CD_3OD) δ 0.89 (t, 3H, $J_{\text{CH}_3,\text{CH}_2} = 7.3$ Hz, $\text{CH}_2\text{-CH}_3$), 1.26–1.37 (m, 8H, $\text{CH}_2\text{-(CH}_2)_4\text{-CH}_3$), 1.57–1.65 (m, 2H, $\text{CH}_2\text{-CH}_2\text{-(CH}_2)_4$), 2.34 (t, 2H, $J = 7.4$ Hz, $\text{CO-CH}_2\text{-CH}_2$), 3.44–3.51 (m, 2H, H-2, H-3) 3.50 (s, 3H, O-CH_3), 3.71 (ddd, 1H, $J_{\text{H-5,H-6}} = 7.3$ Hz, $J_{\text{H-5,H-6}'} = 5.0$ Hz, $J_{\text{H-4,H-5}} = 1.0$ Hz, H-5), 3.79 (d, 1H, $J_{\text{H-3,H-4}} = 2.0$ Hz, H-4), 4.12 (d, 1H, $J_{\text{H-1,H-2}} = 7.5$ Hz, H-1), 4.21 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.4$ Hz, $J_{\text{H-5,H-6}} = 5.0$ Hz, H-6), 4.30 (dd, 1H, $J_{\text{H-6,H-6}'} = 11.3$ Hz, $J_{\text{H-5,H-6}'} = 7.4$ Hz, H-6'), ^{13}C NMR (100 MHz, CD_3OD) δ 14.4 ($-\text{CH}_2\text{-CH}_3$), 23.7, 26.1, 30.1, 30.2, 32.8 (5 s, $-(\text{CH}_2)_5\text{-CH}_3$), 35.0 (CO-CH_2 -), 57.2 (O-CH_3), 64.6 (C-6), 70.3 (C-4), 72.3 (C-2), 73.9 (C-5), 74.7 (C-3), 105.9 (C-1), 175.3 (CO). HRMS (ESI) m/z for $\text{C}_{15}\text{H}_{28}\text{O}_7\text{Na}^+$ $[\text{M} + \text{Na}]^+$ calcd 343.1733, found 343.1743.

3.4. Physicochemical Characterizations

3.4.1. Solubility

Water solubility reflects the ability of a particular molecule to dissolve and its interaction with water. To characterize the water solubility of most amphiphile molecules, especially the ionic surfactants, the Krafft point (T_k) is often used, referring to the critical temperature above which micelles form or the melting point of the hydrated solid surfactant. Two general approaches were used to measure the T_k s of the molecules investigated herein: (1) a visual examination of the dissolution phenomenon for binary molecule/water systems, and (2) a quantitative measurement by using differential scanning calorimetry (DSC) analysis. The visual observation procedure is detailed below. We start with adding the molecule into 5–50 mL water, in a concentration range between 10^{-2} mM and 10^3 mM (corresponding to a range of a surfactant mass concentration well below 20%), keeping the suspension under stirring for a certain time at room temperature (RT). Consequently, molecules can be classified into two major types: those soluble in water at RT and others insoluble at RT, with the solubility limit being called S (expressed in mM). The aspect of the sample, either transparent or turbid, is determined by direct visual examination. T_k determination with DSC is particularly applicable in the case of molecules insoluble at RT, for which a concentrated binary molecule/water mixture with a mass fraction of 20% of each molecule ($C \gg \text{CMC}$ in most cases) was prepared at 65°C under stirring for 30 min to obtain a homogeneous system. These pre-homogenized aqueous systems were cooled and conserved in a refrigerator at 4°C for a suitable period (>7 days), during which equilibrated hydrated crystals formed and precipitated. Then, for each molecule, about a 200–400 mg crystal sample was taken and put into the calorimeter cell. The first equilibrium phase was programmed at 4°C for 1 h, followed by a slow heating stage from 4°C to 65°C at a rate of $0.5^\circ\text{C}/\text{min}$. T_k was determined based on the location of the endothermic peak on the DSC curve, if any. All agitations were performed using a standard magnetic hotplate stirrer, MR 3001K (Heidolph Instruments, Schwabach, Germany). DSC curves were obtained from a commercial μDSC7 evo calorimeter (SETARAM Instrumentation, Caluire-et-Cuire, France). T_k s are reported as the averages of at least two measurements as well as their standard deviations.

3.4.2. Surface Activity

The surface activity reflects the amphiphiles' adsorption behavior on the air–aqueous solution interface. The surface tension (γ) was obtained using the Wilhelmy plate method with a K100 Processor Tensiometer (KRÜSS, Hamburg, Germany). Both the critical micellar concentration (CMC) and equilibrium surface tension at the CMC (γ_{cmc}) were derived from the typical plot of surface tension (γ) against the logarithm of concentration (Log C). The surface excess (Γ_{max}) and minimal area per molecule at the air–aqueous solution interface (A_{min}) at different temperatures were calculated from the Gibbs adsorption equation [3,46]:

$$\Gamma_{\text{max}} = -\frac{1}{2.303RT} \left(\frac{d\gamma}{d\log c} \right) \quad (2)$$

and

$$A_{\text{min}} = \frac{10^{20}}{N_A \Gamma_{\text{max}}} \quad (3)$$

where R is the ideal gas constant, $8.31 \text{ J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$; T is the measuring temperature, K; γ is the surface tension, N/m; c is the concentration, $\text{mol}\cdot\text{L}^{-1}$; A_{min} is the minimal area per molecule, \AA^2 , and N_A is Avogadro's number, $6.022\cdot 10^{23} \text{ mol}^{-1}$.

For each molecule, several solutions of a wide range of concentrations were prepared. Before measuring, each solution was finely homogenized with the help of a magnetic stirring bar. The stirring time was fixed at 30 min in all cases. For molecules with a T_k below RT, the preparation was performed at environmental conditions, whereas for molecules exhibiting a T_k higher than 25°C , the aqueous mixtures were firstly heated and

agitated at elevated temperatures (25–50 °C), depending on the corresponding Tk. Then, measurements were performed once for each concentration, at the solution preparation's temperature, with an uncertainty of ± 1 °C.

4. Conclusions

Two series of methyl D-glycopyranoside-based esters, with different orientations of the ester group, were compared in terms of solubility and surface adsorption properties. This study, combined with a comparison of four well-studied commercial octyl glycosides (C8 α Glc, C8 β Glc, C8 α Gal, and C8 β Gal), allows for a careful exploration of how structural features affect their physicochemical properties.

Commercial octyl D-glycosides presented high solubility limits although their Krafft points were strongly influenced by the head group configuration. Only C8 α Glc and C8 β Gal were found to have Tks above room temperature. Specific structural features causing differences in solubility were proposed for these molecules. For sugar esters families, the ester group orientation had an obvious impact on the dissolution behavior of the compounds.

With respect to the surface tension measurements, we observed, for both series of ester surfactants, a clear linear decrease in the log CMC with the alkyl chain length, in line with well-established surfactant science research. In addition, we could also evidence a slight decrease in γ_{cmc} and A_{min} with alkyl chain lengths, consistent with literature-based expectations for sugar-based surfactants. By comparing the commercial octyl D-glycosides and all the synthesized C7/C8 esters, we found that the latter exhibits a noticeably lower CMC. At a constant alkyl chain length, the various polar heads studied in this work did not seem to impact γ_{cmc} significantly.

Overall, this work offers a refined understanding of the various factors involved in the physicochemical properties of sugar-based amphiphiles. While being able to predict structure-amphiphilic property trends should allow the design of greener surfactants in a smarter way, it is obvious that predicting the solubility properties of new amphiphilic molecules is tricky and challenging, as sugar-based surfactant solubility arises from a delicate balance between three key factors—their crystal stability, hydrophilicity, and self-assembling tendency. Further research on the structure–property relationships of sugar-based surfactants are still needed to contribute to community knowledge and to provide significant hope regarding the replacement of petroleum-based substances.

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/molecules29102338/s1>, ^1H and ^{13}C NMR spectra of final compounds (8–12).

Author Contributions: A.D., A.W., G.P. and I.P. conceived and designed the experiments; H.L. and B.B. performed the experiments; all authors analyzed the data; T.G. notably contributed to the structure–property relationships; G.P. and A.D. were the primary writers of the manuscript; all authors proofread the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by a grant awarded to B.B. and PhD funding for T.G., provided as part of the French Government's Investments for the Future, under the reference ANR-001-01.

Institutional Review Board Statement: Not Applicable.

Informed Consent Statement: Not Applicable.

Data Availability Statement: The original contributions presented in the study are included in the article/Supplementary Materials, further inquiries can be directed to the corresponding author/s.

Acknowledgments: H.L. acknowledges the Chinese Scholarship Council for the financial support of her PhD grant. This work was performed in partnership with SAS PIVERT, within the framework of the French Institute for the Energy Transition (Institut pour la Transition Énergétique (ITE) P.I.V.E.R.T. (www.institutpivert.com)).

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Von Rybinski, W.; Hill, K. Alkyl polyglycosides—Properties and applications of a new class of surfactants. *Angew. Chem. Int. Ed.* **1998**, *37*, 1328–1345. [[CrossRef](#)]
2. Hill, K.; Rhode, O. Sugar-based surfactants for consumer products and technical applications. *Fett-Lipid* **1999**, *101*, 25–33. [[CrossRef](#)]
3. Ruiz, C.C. *Sugar-Based Surfactants: Fundamentals and Applications*; CRC Press: Boca Raton, FL, USA, 2008; pp. 245–306.
4. Rojas, O.J.; Stubenrauch, C.; Lucia, L.A.; Habibi, Y. Interfacial properties of sugar-based surfactants. In *Biobased Surfactants and Detergents: Synthesis, Properties, and Applications*; Hayes, D., Kitamoto, D., Solaiman, D., Ashby, R., Eds.; AOCS Press: Urbana, IL, USA, 2009; p. 457.
5. Pal, A.; Mondal, M.H.; Adhikari, A.; Bhattarai, A.; Saha, B. Scientific information about sugar-based emulsifiers: A comprehensive review. *RSC Adv.* **2021**, *11*, 33004–33016. [[CrossRef](#)] [[PubMed](#)]
6. Gaudin, T.; Lu, H.; Fayet, G.; Berthaud-Drelich, A.; Rotureau, P.; Pourceau, G.; Wadouachi, A.; Van Hecke, E.; Nesterenko, A.; Pezron, I. Impact of the chemical structure on amphiphilic properties of sugar-based surfactants: A literature overview. *Adv. Colloid Interface Sci.* **2019**, *270*, 87–100. [[CrossRef](#)] [[PubMed](#)]
7. Razafindralambo, H.; Blecker, C.; Paquot, M. Screening of basic properties of amphiphilic molecular structures for colloidal system formation and stability. In *Amphiphiles: Molecular Assembly and Applications*; American Chemical Society: Washington, DC, USA, 2011; Volume 1070, Chapter 4; pp. 53–66.
8. Razafindralambo, H.; Blecker, C.; Paquot, M. 8. Carbohydrate-based surfactants: Structure-activity relationships. In *Advances in Chemical Engineering*; Nawaz, Z., Ed.; InTech: Rijeka, Croatia, 2012.
9. Piispanen, P.S.; Persson, M.; Claesson, P.; Norin, T. Surface properties of surfactants derived from natural products. Part 1: Syntheses and structure/property relationships—Solubility and emulsification. *J. Surf. Det.* **2004**, *7*, 147–159. [[CrossRef](#)]
10. Lu, B.; Vayssade, M.; Miao, Y.; Chagnault, V.; Grand, E.; Wadouachi, A.; Postel, D.; Drelich, A.; Egles, C.; Pezron, I. Physico-chemical properties and cytotoxic effects of sugar-based surfactants: Impact of structural variations. *Colloids Surf. B* **2016**, *145*, 79–86. [[CrossRef](#)]
11. Baker, I.J.A.; Matthews, B.; Soares, H.; Krodkiewska, I.; Furlong, D.N.; Grieser, F.; Drummond, C.J. Sugar fatty acid ester surfactants: Structure and ultimate aerobic biodegradability. *J. Surf. Det.* **2000**, *3*, 1–11. [[CrossRef](#)]
12. Becerra, N.; Toro, C.; Zanocco, A.L.; Lemp, E.; Günther, G. Characterization of micelles formed by sucrose 6-O-monoesters. *Colloids Surf. A* **2008**, *327*, 134–139. [[CrossRef](#)]
13. Cook, A.G.; Wardell, J.L.; Imrie, C.T. Carbohydrate liquid crystals: Synthesis and characterisation of the methyl-6-O-(n-acyl)- α -D-glucopyranosides. *Chem. Phys. Lipids* **2011**, *164*, 118–124. [[CrossRef](#)]
14. Razafindralambo, H.; Richel, A.; Wathelet, B.; Blecker, C.; Wathelet, J.P.; Brasseur, R.; Lins, L.; Minones, J.; Paquot, M. Monolayer properties of uronic acid bicatenary derivatives at the air-water interface: Effect of hydroxyl group stereochemistry evidenced by experimental and computational approaches. *Phys. Chem. Chem. Phys.* **2011**, *13*, 15291–15298. [[CrossRef](#)]
15. Ducret, A.; Giroux, A.; Trani, M.; Lortie, R. Enzymatic preparation of biosurfactants from sugars or sugar alcohols and fatty acids in organic media under reduced pressure. *Biotechnol. Bioeng.* **1995**, *48*, 214–221. [[CrossRef](#)] [[PubMed](#)]
16. Skagerlind, P.; Larsson, K.; Barfoed, M.; Hult, K. Glucoside ester synthesis in microemulsions catalyzed by *Candida antarctica* component B lipase. *J. Am. Oil Chem. Soc.* **1997**, *74*, 39–42. [[CrossRef](#)]
17. Blecker, C.; Piccicuto, S.; Lognay, G.; Deroanne, C.; Marlier, M.; Paquot, M. Enzymatically prepared n-alkyl esters of glucuronic acid: The effect of hydrophobic chain length on surface properties. *J. Colloid Int. Sci.* **2002**, *247*, 424–428. [[CrossRef](#)] [[PubMed](#)]
18. Moreau, B.; Lognay, G.C.; Blecker, C.; Brohée, J.C.; Chéry, F.; Rollin, P.; Paquot, M.; Marlier, M. Synthesis of novel D-glucuronic acid fatty esters using *Candida antarctica* lipase in tert-butanol. *Biotechnol. Lett.* **2004**, *26*, 419–424. [[CrossRef](#)] [[PubMed](#)]
19. Gouéth, P.Y.; Gogalis, P.; Bikanga, R.; Godé, P.; Postel, D.; Ronco, G.; Villa, P. Synthesis of Monoesters as Surfactants and Drugs from D-Glucose. *J. Carbohydr. Chem.* **1994**, *13*, 249–272. [[CrossRef](#)]
20. Otto, R.T.; Bornscheuer, U.T.; Syldatk, C.; Schmid, R.D. Lipase-catalyzed synthesis of arylaliphatic esters of β -D(+)-glucose, n-alkyl- and arylglucosides and characterization of their surfactant properties. *J. Biotechnol.* **1998**, *64*, 231–237. [[CrossRef](#)]
21. Savelli, M.P.; Van Roekeghem, P.; Douillet, O.; Cavé, G.; Godé, P.; Ronco, G.; Villa, P. Effects of tail alkyl chain length (n), head group structure and junction (Z) on amphiphilic properties of 1-Z-R-D,L-xylitol compounds ($R=C_nH_{2n+1}$). *Int. J. Pharm.* **1999**, *182*, 221–236. [[CrossRef](#)] [[PubMed](#)]
22. Garofalakis, G.; Murray, B.S.; Sarney, D.B. Surface Activity and Critical Aggregation Concentration of Pure Sugar Esters with Different Sugar Headgroups. *J. Colloid Interface Sci.* **2000**, *229*, 391–398. [[CrossRef](#)] [[PubMed](#)]
23. Kjellin, U.R.M.; Claesson, P.M.; Vulfson, E.N. Studies of N-Dodecyl lactobionamide, Maltose 6'-O-Dodecanoate, and Octyl- β -glucoside with Surface Tension, Surface Force, and Wetting Techniques. *Langmuir* **2001**, *17*, 1941–1949. [[CrossRef](#)]
24. Soultani, S.; Ognier, S.; Engasser, J.M.; Ghoul, M. Comparative study of some surface-active properties of fructose esters and commercial sucrose esters. *Colloids Surf. A* **2003**, *227*, 35–44. [[CrossRef](#)]
25. Brown, G.M.; Dubreuil, P.; Ichhaporia, F.M.; Desnoyers, J.E. Synthesis and properties of some α -D-alkyl glucosides and mannosides: Apparent molal volumes and solubilization of nitrobenzene in water at 25 °C. *Can. J. Chem.* **1970**, *48*, 2525–2531. [[CrossRef](#)]
26. Nilsson, F.; Söderman, O.; Johansson, I. Physical–Chemical Properties of the n-Octyl β -D-Glucoside/Water System. A Phase Diagram, Self-Diffusion NMR, and SAXS Study. *Langmuir* **1996**, *12*, 902–908. [[CrossRef](#)]

27. Dupuy, C.; Auvray, X.; Petipas, C.; Rico-Lattes, I.; Lattes, A. Anomeric effects on the structure of micelles of alkyl maltosides in water. *Langmuir* **1997**, *13*, 3965–3967. [CrossRef]
28. Fukada, K.; Kawasaki, M.; Seimiya, T.; Abe, Y.; Fujiwara, M.; Ohbu, K. Stereochemical aspects of micellar properties of esterified glucoside surfactants in water: Apparent molar volume, adiabatic compressibility, and aggregation number. *Colloid Polym. Sci.* **2000**, *278*, 576–580. [CrossRef]
29. Lu, H.; Drelich, A.; Omri, M.; Pezron, I.; Wadouachi, A.; Pourceau, G. Catalytic synthesis of a new series of alkyl uronates and evaluation of their physicochemical properties. *Molecules* **2016**, *21*, 1301. [CrossRef] [PubMed]
30. Lu, H.; Pezron, I.; Gaudin, T.; Drelich, A. Non-equilibrium micelles formed by sugar-based surfactants under their Krafft temperature. *Colloids Surf. A* **2018**, *540*, 167–176. [CrossRef]
31. Hato, M. Synthetic glycolipid/water systems. *Curr. Opin. Colloid Interface Sci.* **2001**, *6*, 268–276. [CrossRef]
32. Matsumura, S.; Imai, K.; Yoshikawa, S.; Kawada, K.; Uchibor, T. Surface activities, biodegradability and antimicrobial properties of n-alkyl glucosides, mannosides and galactosides. *J. Am. Oil Chem. Soc.* **1990**, *67*, 996–1001. [CrossRef]
33. Schmidt-Lassen, J.; Lindhorst, T.K. Exploring the meaning of sugar configuration in a supramolecular environment: Comparison of six octyl glycoside micelles by ITC and NMR spectroscopy. *MedChemComm* **2014**, *5*, 1218–1226. [CrossRef]
34. Khan, A.A.; Chee, S.H.; McLaughlin, R.J.; Harper, J.L.; Kamena, F.; Timmer, M.S.M.; Stocker, B.L. Long-Chain Lipids Are Required for the Innate Immune Recognition of Trehalose Diesters by Macrophages. *ChemBioChem* **2011**, *12*, 2572–2576. [CrossRef]
35. Kotena, Z.M.; Behjatmanesh-Ardakani, R.; Hashim, R. AIM and NBO analyses on hydrogen bonds formation in sugar-based surfactants (α/β -D-mannose and n-octyl- α/β -D-mannopyranoside): A density functional theory study. *Liq. Cryst.* **2014**, *41*, 784–792. [CrossRef]
36. Straathof, A.J.J.; Van Bekkum, H.; Kieboom, A.P.G. Solid State and Solution Properties of Octyl D-Glucopyranosides. *Starch* **1988**, *40*, 438–440. [CrossRef]
37. Sakya, P.; Seddon, J.M. Thermotropic and lyotropic phase behaviour of monoalkyl glycosides. *Liq. Cryst.* **1997**, *23*, 409–424. [CrossRef]
38. Boyd, B.J.; Drummond, C.J.; Krodziewska, I.; Grieser, F. How Chain Length, Headgroup Polymerization, and Anomeric Configuration Govern the Thermotropic and Lyotropic Liquid Crystalline Phase Behavior and the Air–Water Interfacial Adsorption of Glucose-Based Surfactants. *Langmuir* **2000**, *16*, 7359–7367. [CrossRef]
39. Jmol Development Team. Jmol. 2016. Available online: <http://jmol.sourceforge.net/> (accessed on 22 April 2024).
40. Rosen, M.J.; Kunjappu, J.T. *Surfactants and Interfacial Phenomena*, 4th ed.; Wiley: Hoboken, NJ, USA, 2006; ISBN 978-0-470-54194-4.
41. Capalbi, A.; Gente, G.; La Mesa, C. Solution properties of alkyl glucosides, alkyl thioglucosides and alkyl maltosides. *Colloids Surf. A* **2004**, *246*, 99–108. [CrossRef]
42. Lainez, A.; Del Burgo, P.; Junquera, E.; Aicart, E. Mixed Micelles Formed by n-Octyl- β -D-glucopyranoside and Tetradecyltrimethylammonium Bromide in Aqueous Media. *Langmuir* **2004**, *20*, 5745–5752. [CrossRef] [PubMed]
43. Mańko, D.; Zdziennicka, A.; Jańczuk, B. Thermodynamic properties of adsorption and micellization of n-octyl- β -D-glucopyranoside. *Colloids Surf. B* **2014**, *114*, 170–176. [CrossRef] [PubMed]
44. Silva, F.V.; Goulart, M.; Justino, J.; Neves, A.; Santos, F.; Caio, J.; Lucas, S.; Newton, A.; Sacoto, D.; Barbosa, E.; et al. Alkyl deoxy-arabino-hexopyranosides: Synthesis, surface properties, and biological activities. *Bioorganic Med. Chem.* **2008**, *16*, 4083–4092. [CrossRef]
45. Frindi, M.; Michels, B.; Zana, R. Ultrasonic absorption studies of surfactant exchange between micelles and bulk phase in aqueous micellar solutions of nonionic surfactants with a short alkyl chain. 3. Surfactants with a sugar head group. *J. Phys. Chem.* **1992**, *96*, 8137–8141. [CrossRef]
46. Shinoda, K.; Yamaguchi, T.; Hori, R. The Surface Tension and the Critical Micelle Concentration in Aqueous Solution of β -D-Alkyl Glucosides and their Mixtures. *Bull. Chem. Soc. Jpn.* **1961**, *34*, 237–241. [CrossRef]
47. Sulthana, S.B.; Rao, P.V.C.; Bhat, S.G.T.; Nakano, T.Y.; Sugihara, G.; Rakshit, A.K. Solution Properties of Nonionic Surfactants and Their Mixtures: Polyoxyethylene (10) Alkyl Ether [CnE10] and MEGA-10. *Langmuir* **2000**, *16*, 980–987. [CrossRef]
48. Wang, X.; Yan, F.; Li, Z.; Zhang, L.; Zhao, S.; An, J.; Yu, J. Synthesis and surface properties of several nonionic–anionic surfactants with straight chain alkyl-benzyl hydrophobic group. *Colloids Surf. A* **2007**, *302*, 532–539. [CrossRef]
49. Alami, E.; Beinert, G.; Marie, P.; Zana, R. Alkanediyl- α,ω -bis(dimethylalkylammonium bromide) surfactants. 3. Behavior at the air–water interface. *Langmuir* **1993**, *9*, 1465–1467. [CrossRef]
50. Ueno, M.; Takasawa, Y.; Miyashige, H.; Tabata, Y.; Meguro, K. Effects of alkyl chain length on surface and micellar properties of octaethyleneglycol-n alkyl ethers. *Colloid Polym. Sci.* **1981**, *259*, 761–766. [CrossRef]
51. Zhou, T.; Yang, H.; Xu, X.; Wang, X.; Wang, J.; Dong, G. Synthesis, surface and aggregation properties of nonionic poly(ethylene oxide) gemini surfactants. *Colloids Surf. A* **2008**, *317*, 339–343. [CrossRef]
52. Razafindralambo, H.; Blecker, C.; Mezdour, S.; Deroanne, C.; Crowet, J.M.; Bresseur, R.; Lins, L.; Paquot, M. Impacts of the Carbonyl Group Location of Ester Bond on Interfacial Properties of Sugar-Based Surfactants: Experimental and Computational Evidences. *J. Phys. Chem. B* **2009**, *113*, 8872–8877. [CrossRef] [PubMed]

53. Li, G.-L.; Kung, K.K.-Y.; Wong, M.-K. Gold-catalyzed amide synthesis from aldehydes and amines in aqueous medium. *Chem. Comm.* **2012**, *48*, 4112–4114. [[CrossRef](#)]
54. Wang, H.; Cui, Y.; Zou, R.; Cheng, Z.; Yao, W.; Mao, Y.; Zhang, Y. Synthesis of oligosaccharides using per-O-trimethylsilyl-glycosyl iodides as glycosyl donors. *Carbohydr. Res.* **2016**, *427*, 1–5. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.